www.tntextbooks.in

Flowers: Bracteate, bracteolate (*Sesbania*), pedicellete, complete, bisexual, pentamerous, heterochlamydeous, zygomorphic hypogynous or sometimes perigynous.

Calyx: Sepals 5, green, synsepalous, more or less united in a tube and persistant, valvate or imbricate, odd sepal is anterior in position.

Corolla: Petals 5, apopetalous, unequal and **papilionaceous**, **vexillary** or **descendingly imbricate** aestivation all petals have claw at the base. The outer most petal is large called **standard petal** or **vexillum**, Lateral 2 petals are lanceolate and curved. They are called **wing petals** or **alae**. Anterior two petals are partly fused and are called **keel petals** or **carina** which encloses the stamens and pistil.

Androecium: Stamens 10, diadelphous, usually 9+1 (*Clitoria ternatea*). The odd stamen is posterior in position. In *Aeschynomene aspera*, the stamens are fused to form two bundles each containing five stamens (5)+5. Stamens are **monadelphous** and **dimorphic** ie. 5 stamens have longer filaments and other 5 stamens have shorter filaments thus the stamens are found at two levels and the shape of anthers also varies in (*Crotalaria verrucosa)*. (5 anthers are long and lanceolate, and the other 5 anthers are short and blunt). Anthers are dithecous, basifixed and dehiscing longitudinally.

Gynoecium: Monocarpellary, unilocular, ovary superior, with two alternating rows of ovules on marginal placentation. Style simple and bent, stigma flattened or feathery.

Fruit: The characteristic fruit of Fabaceae is a legume (*Pisum sativum)*, sometimes indehiscent and rarely a lomentum (*Desmodium)*.

In *Arachis hypogea* the fruit is ***geocarpic*** (fruits develops and matures from underground). After fertilization the stipe of the ovary becomes meristematic and grows down into the soil. This ovary gets buried into the soil and develops into fruit.

Seed: Endospermic or non-endospermic (*Pisum sativum*), mostly **reniformed**.

Botanical description of *Clitoria ternatea* (Sangu pushpam)

Habit: Twining climber

Root: Branched tap root system having nodules.

Stem: Aerial, weak stem and a twiner

Leaf: Imparipinnately compound, alternate, stipulate showing reticulate venation. Leaflets are stipellate. Petiolate and stipels are pulvinated.

Inflorescence: Solitary and axillary

Flower: Bracteate, bracteolate, bracteoles usually large, pedicellate, heterochlamydeous, complete, bisexual, pentamerous, zygomorphic and hypogynous.

Calyx: Sepals 5, synsepalous, green showing valvate aestivation. Odd sepal is anterior in position.

Corolla: Petals 5, white or blue apopetalous, irregular papilionaceous corolla showing descendingly imbricate aestivation.

Androecium: Stamens 10, diadelphous (9)+1 nine stamens fused to form a bundle and the tenth stamen is free. Anthers