www.tntextbooks.in

in species of *Smilax* it is reticulate. Leaves are usually exstipulate, but in *Smilax*, two tendrils arise from the base of the leaf, which are considered modified stipules.

Inflorescence: Flowers are usually borne in simple or branched racemes (*Asphodelus*) spikes in *Aloe*, huge terminal panicle in *Yucca*, solitary and axillary in *Gloriosa*, solitary and terminal in *Tulipa.*

Flowers: Flowers are often showy, pedicellate, bracteate, usually ebracteolate except *Dianella* and *Lilium*, bisexual, actinomorphic, trimerous, hypogynous, rarely unisexual (*Smilax*) and are dioecious, rarely tetramerous (*Maianthemum*), slightly zygomorphic (*Lilium*) and hypogynous.

Perianth: Tepals 6 biseriate arranged in two whorls of 3 each, apotepalous or rarely syntepalous as in *Aloe.* Usually petaloid or sometimes sepaloid, odd tepal of the outer whorl is anterior in position, valvate or imbricate, tepals more than six in *Paris quadrifolia*.

Androecium: Stamens 6, arranged in 2 whorls of 3 each: rarely stamens are 3 (*Ruscus*),4 in *Maianthemum,* or up to 12, apostamenous, opposite to the tepals, sometimes epitepalous; filaments distinct or connate, anthers dithecous, basifixed or versatile, extrose, or intrese, dehiscing usually by vertical slit and sometimes by terminal pores; rarely **synstamenous** (*Ruscus*).

Gynoecium: Tricarpallary, syncarpous, the odd carpel usually anterior, ovary superior, trilocular, with 2 rows of numerous ovules on axile placextation; rarely unilocular with parietal placentation, style usually one; stigmas 1 or 3; rarely the ovary is inferior (*Haemodorum*), nectar – secreting **septal glands** are present in the ovary.

Fruit: Fruit usually a septicidal or loculicidal capsule or a berry as in *Asparagus* & *Smilax.*

Botanical description of *Allium cepa*

(In APG classification, *Allium cepa* is placed under the family Amaryllidaceae)

Habit: Perennial herb with bulb.

Root: Fibrous adventitious root system

Stem: Underground bulb

Leaf: a cluster of radical leaves emerges from the underground bulb, cylindrical and fleshy having sheathy leaf bases with parallel venation.

Inflorescence: Scapigerous i.e. the inflorescence axis (peduncle) arising from the ground bearing a cluster of flowers at its apex. Pedicels are of equal length, arising from the apex of the peduncle which brings all flowers at the same level.

Flower: Small, white, bracteate, ebrcteolate, pedicellate, complete, trimerous, actinomorphic and hypogynous. Flowers are protandrous.

Perianth: Tepals 6, white,arranged in two whorls of three each, syntepalous showing valvate aestivation.

Androecium: Stamens 6, arranged in two whorls of three each, epitepalous, apostamenous /free and opposite to tepals. Anthers dithecous, basifixed, introse, and dehiscing longitudinally.

Gynoecium: Tricarpellary and syncarpous. Ovary superior, trilocular with two ovules in each locule on axile placentation. Style simple, slender with simple stigma.

Fruit: A loculicidal capsule.

Seed: Endospermous

Floral Formula:

$$\text{Br.,Ebrl.,}\oplus,\male\female,P_{(3+3)}+A_{3+3},\underline{G}_{(3)}$$