www.tntextbooks.in

Competitive Exam Questions

Unit -4 – Plant Anatomy

1. The balloon – shaped structures called tyloses (NEET II – 2016)
   a. originate in the lumen of vessels
   b. characterise the sap wood
   c. are extensions of xylem parenchyma cells into vessels
   d. are linked to the ascent of sap through xylem vessels

2. Cortex is the region found between (NEET II – 2016)
   a. epidermis and stele
   b. pericycle and endodermis
   c. endodermis and pith
   d. endodermis and vascular bundle

3. Read I – IV and find the correct order of components from outer side to inner side in a woody dicot stem (CBSE -AIPMT – 2015)
   (I) secondary Cortex (II) wood
   (III) secondary phloem (IV) phellem
   a. III, IV, II and I b. I, II, IV and III
   c. IV, I, III and II d. IV, III, I and II

4. You are given a fairly old piece of a dicot stem and a dicot root. Which of the following anatomical structures will you use to distinguish between the two? (CBSE -AIPMT 2014)
   a. secondary xylem
   b. secondary phloem
   c. protoxylem
   d. cortical cells

5. Heart wood differs from sapwood in (CBSE -AIPMT 2010)
   a. the presence of rays and fibres
   b. the absence of vessels and parenchyma
   c. having dead and non-conducting elements
   d. being susceptible to hosts and pathogens

6. The annular and spirally thickened conducting elements generally develop in the protoxylem when the root or stem is (CBSE -AIPMT 2009)
   a. maturing b. elongating
   c. widening d. differentiating

7. Anatomically fairly old dicotyledonous root is distinguished from the dicotyledonous stem by the (CBSE- AIPMT 2009)
   a. absence of secondary xylem
   b. absence of secondary phloem
   c. presence of cortex
   d. position of protoxylem

8. In barley stem, vascular bundles are (CBSE -AIPMT 2009)
   a. open and scattered
   b. closed and scattered
   c. open and in a ring
   d. closed and radial

9. Palisade parenchyma is absent in the leaves of (CBSE- AIPMT 2009)
   a. sorghum b. mustard
   c. soyabean d. gram

10. Sugarcane plant has (AIIMS 2009)
    a. reticulate venation
    b. capsular fruits
    c. pentamerous flowers
    d. dump-bell shaped guard cells

11. Vascular tissues in flowering plants develop from (CBSE- AIPMT 2008 & JIPMER 2012)
    a. phellogen **b. plerome**
    c. periblem d. dermatogen

12. The length of different internodes in a culm of sugarcane is variable because of (CBSE -AIPMT 2008)
    a. short apical meristem
    b. position of axillary buds