Placental Lactogen (hPL), oestrogens and progesterone which are essential for a normal pregnancy. A hormone called **relaxin** is also secreted during the later phase of pregnancy which helps in relaxation of the pelvic ligaments at the time of parturition. It should be noted that hCG, hPL and relaxin are produced only during pregnancy. In addition, during pregnancy the level of other hormones like oestrogen and progesterone, cortisol, prolactin, thyroxine, etc., is increased several folds in the maternal blood. These hormones are essential for supporting foetal growth.



A female uterus is normally about 3 inches long and 2 inches wide but can expand 20 times during pregnancy.

2.6 Parturition and lactation

Parturition is the completion of pregnancy and giving birth to the baby. The series of events that expels the infant from the uterus is collectively called "labour". Throughout pregnancy the uterus undergoes periodic episodes of weak and strong contractions. These contractions called **Braxton-Hick's contractions** lead to false labour. As the pregnancy progresses, increase in the oestrogen concentration promotes uterine contractions. These uterine contractions facilitate moulding of the foetus and downward movement of the foetus. The descent of the foetus causes dilation of cervix of the uterus and vaginal canal resulting in a **neurohumoral reflex** called **Foetal ejection reflex** or **Ferguson reflex**. This initiates the secretion of oxytocin from the neurohypophysis which in turn brings about the powerful contraction of the uterine muscles and leads to the expulsion of the baby through the birth canal. This sequence of events is called as **parturition** or **childbirth**.

Relaxin is a hormone secreted by the placenta and also found in the corpus luteum. It promotes parturition by relaxing the pelvic joints and by dilatation of the cervix with continued powerful contractions. The amnion ruptures and the amniotic fluid flows out through the vagina, followed by the foetus. The placenta along with the remains of the umbilical cord called **"after birth"** is expelled out after delivery.

Lactation is the production of milk by mammary glands. The mammary glands show changes during every menstrual cycle, during pregnancy and lactation. Increased level of oestrogens, progesterone and **human Placental Lactogen (hPL)** towards the end of pregnancy stimulate the hypothalamus towards prolactin – releasing factors. The anterior pituitary responds by secreting **prolactin** which plays a major role in lactogenesis.

Oxytocin causes the "**Let-Down" reflex**- the actual ejection of milk from the alveoli of the mammary glands. During lactation, oxytocin also stimulates the recently emptied uterus to contract, helping it to return to pre - pregnancy size.

Colostrum

Colostrum, a nutrient rich fluid produced by the human female immediately after giving birth, is loaded with immune, growth and tissue repair factors. It acts as a natural antimicrobial agent to actively stimulate the maturation of the infant's immune system. No artificial feed can substitute the first milk, with all its natural benefits and therefore should be definitely fed to the baby after birth.

The mammary glands secrete a yellowish fluid called **colostrum** during the initial few days after parturition. It has less lactose

XII Std Biology-Zoology Chapter-2 EM.indd 24 23/12/2022 11:08:36