CASE STUDY

Approximately half of unintended pregnancies are due to contraceptive failure, largely owing to inconsistent or incorrect use of contraceptive methods. The effectiveness of **long-acting reversible contraception** (intrauterine devices and contraceptive implants) is superior to that of contraceptive pills, patch, or ring and is not altered in adolescents and young women. Educating young women about the usage of intrauterine devices and contraceptive implants would dramatically reduce the number of unintended pregnancies among women seeking family planning.

pregnancy and generally has no impact on a women's fertility. Abortion during the second trimester is more risky as the foetus becomes intimately associated with the maternal tissue. Government of India legalized MTP in 1971 for medical necessity and social consequences with certain restrictions like sex discrimination and illegal female foeticides to avoid its misuse. MTP performed illegally by unqualified quacks is unsafe and could be fatal. MTP of the first conception may have serious psychological consequences

3.6. Sexually Transmitted Diseases (STD)

HG1BS

Sexually transmitted diseases (STD) or Venereal diseases (VD) or Reproductive tract infections (RTI) are called as Sexually transmitted infections (STI). Normally STI are transmitted from person to person during intimate sexual contact with an infected partner. Infections like **Hepatitis-B** and **HIV** are transmitted sexually as well as by sharing of infusion needles, surgical instruments, etc with infected people, blood transfusion or from infected mother to baby. People in the age of 15 to 24 years are prone to these infections. The bacterial STI are gonorrhoea, syphilis, chancroid, chlamydiasis and lymphogranuloma venereum. The viral STI are genital herpes, genital warts, Hepatitis-B and AIDS**.** Trichomoniasis is a protozoan STI, and candidiasis is a fungal STI**.** STI caused by bacteria, fungi and protozoa or parasites, can be treated with antibiotics or other medicines, whereas STI caused by virus cannot be treated but the symptoms can be controlled by antiviral medications. Latex condoms usage greatly reduces the risk, but does not completely eliminate the risk of transmission of STI.

Prevention of STDs

a. Avoid sex with unknown partner/ multiple partners.

b. Use condoms.

c. In case of doubt, consult a doctor for diagnosis and get complete treatment.

DO YOU KNOW?

According to World Health Organization (WHO), 2017 more than one million people globally acquires a sexually transmitted infection every day. India has the third largest HIV epidemic in the world, with 2.1 million people living with HIV.

DO YOU KNOW?

TNHSP (Tamilnadu health systems project), a unit of the Health and family welfare department of the Government of Tamilnadu does free screening for cervical and breast cancer.

XII Std Biology-Zoology Chapter-3 EM.indd 36 23/12/2022 11:17:31