Table 7.4 Innate immunity- types and mechanisms

Type of innate immunity	Mechanism
1. Anatomical barriers	
Skin	Prevents the entry of microbes. Its acidic environment (pH 3-5) retards the growth of microbes.
Mucus membrane	Mucus entraps foreign microorganisms and competes with microbes for attachment.
2. Physiological barriers	
Temperature	Normal body temperature inhibits the growth of pathogens. Fever also inhibits the growth of pathogens.
Low pH	Acidity of gastric secretions (HCl) kills most ingested microbes.
Chemical mediators	Lysozyme acts as antibacterial agent and cleaves the bacterial cell wall. Interferons induce antiviral state in the uninfected cells. Complementary substances produced from leucocytes lyse the pathogenic microbes or facilitate phagocytosis.
3. Phagocytic barriers	Specialized cells (Monocytes, neutrophils, tissue macrophages) phagocytose, and digest whole microorganisms.
4. Inflammatory barriers	Tissue damage and infection induce leakage of vascular fluid, containing chemotactic signals like serotonin, histamine and prostaglandins. They influx the phagocytic cells into the affected area. This phenomenon is called diapedesis.

7.3.2 Acquired immunity

The immunity that an individual acquires after birth is known as **acquired immunity**. It is the body's resistance to a specific pathogen.

The unique features of acquired immunity are antigenic specificity, diversity, recognition of self and non-self and immunological memory.

Components of acquired immunity

Acquired immunity has two components – **cell mediated immunity (CMI) and antibody mediated immunity or humoral immunity.**

1. Cell mediated immunity

When pathogens are destroyed by cells without producing antibodies, then it is known as cell mediated immune response or cell mediated immunity. This is brought about by T cells, macrophages and natural killer cells.

2. Antibody mediated immunity or humoral immunity

When pathogens are destroyed by the production of antibodies, then it is known as antibody mediated or humoral immunity. This is brought about by B cells with the help of antigen presenting cells and T helper cells. Antibody production is the characteristic feature of **vertebrates** only.

Types of acquired immunity

Acquired immunity may be **active immunity** or **passive immunity (Table 7.5)**.

XII Std Biology-Zoology Chapter-7 EM.indd 117 23/12/2022 12:39:56