19. Conductivity of a saturated solution of a sparingly soluble salt AB (1:1 electrolyte) at 298K is 1.85×10^{-5} S m^{-1}. Solubility product of the salt AB at 298K $\left(\Lambda^{\circ}_{m}\right)_{AB} = 14\times10^{-3}$ S m^{2} mol^{-1}.

a) 5.7×10^{-12} b) 1.32×10^{-12} c) 7.5×10^{-12} d) 1.74×10^{-12}

20. In the electrochemical cell: $Zn|ZnSO_4\ (0.01M)\|CuSO_4(1.0M)|Cu$, the emf of this Daniel cell is E_1. When the concentration of $ZnSO_4$ is changed to 1.0M and that $CuSO_4$ changed to 0.01M, the emf changes to E_2. From the above, which one is the relationship between E_1 and E_2?

a) $E_1 < E_2$ b) $E_1 > E_2$ c) $E_2 \geq E_1$ d) $E_1 = E_2$

21. Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below:

$$BrO_4^- \xrightarrow{1.82V} BrO_3^- \xrightarrow{1.5V} HBrO \xrightarrow{1.595V} Br_2 \xrightarrow{1.0652V} Br^-$$

Then the species undergoing disproportionation is

a) Br_2 b) BrO_4^- c) BrO_3^- d) HBrO

22. For the cell reaction

$2Fe^{3+}(aq) + 2I^-(aq) \rightarrow 2Fe^{2+}(aq) + I_2(aq)$

$E^{\circ}_{cell} =$ 0.24V at 298K. The standard Gibbs energy (Δ, G°) of the cell reactions is :

a) -46.32 KJ mol^{-1} b) -23.16 KJ mol^{-1} c) 46.32 KJ mol^{-1} d) 23.16 KJ mol^{-1}

23. A certain current liberated 0.504gm of hydrogen in 2 hours. How many grams of copper can be liberated by the same current flowing for the same time through copper sulphate solution

a) 31.75 b) 15.8 c) 7.5 d) 63.5

24. A gas X at 1 atm is bubbled through a solution containing a mixture of $1MY^-$ and $1MZ^-$ at 25°C. If the reduction potential of Z>Y>X, then

a) Y will oxidize X and not Z b) Y will oxidize Z and not X

d) Y will oxidize both X and Z d) Y will reduce both X and Z

25. Cell equation : $A^{2+} + 2B^- \rightarrow A + 2B$

$A^{2+} + 2e^- \rightarrow A \quad E^{\circ} = +0.34V$ and $\log_{10}K = 15.6$ at 300K for cell reactions find E° for $B^+ + e^- \rightarrow B$ (AIIMS – 2018)

a) 0.80 b) 1.26 c) -0.54 d) -10.94

Short Answer Questions

1. Define anode and cathode
2. Why does conductivity of a solution decrease on dilution of the solution
3. State Kohlrausch Law. How is it useful to determine the molar conductivity of weak electrolyte at infinite dilution.

XII U9 Electro Chemistry.indd 65 16-12-2022 19:37:47