
enter and could not escape because they are trapped by the hairs. As soon as the anthers of the flower ripe, the hairs wither and flower bents down. The flies escape with pollen and enter another flower where it dusts pollen on the stigma bringing out pollination.

Significance of Pollination

- Pollination is a pre-requisite for the process of fertilisation. Fertilisation helps in the formation of fruits and seeds.
- It brings the male and female gametes closer for the process of fertilisation.
- Cross-pollination introduces variations in plants due to the mixing up of different genes. These variations help the plants to adapt to the environment and results in speciation.

1.5 Fertilization



The fusion of male and female gamete is called **fertilization**. **Double fertilization** is seen in angiosperms.

Events of fertilization

The stages involved in double fertilization are:- germination of pollen to form pollen tube in the stigma; growth of pollen tube in the style; direction of pollen tube towards the micropyle of the ovule; entry of the pollen tube into one of the synergids of the embryo sac, discharge of male gametes; syngamy and triple fusion. The events from pollen deposition on the stigma to the entry of pollen tube in to the ovule is called **pollen-pistil interaction.** It is a dynamic process which involves recognition of pollen and to promote or inhibit its germination and growth.

Pollen on the stigma

In nature, a variety of pollens fall on the receptive stigma, but all of them do not germinate and bring out fertilization. The receptive surface of the stigma receives the pollen. If the pollen is compatible with the stigma it germinates to form a tube. This is facilitated by the stigmatic fluid in **wet stigma** and pellicle in **dry stigma**. These two also decide the incompatibility and compatibility of the pollen through **recognition-rejection protein reaction** between the pollen and stigma surface. Sexual incompatibility may exist between different species (interspecific) or between members of the same species (intraspecific).The latter is called self-incompatibility. The first visible change in the pollen, soon after it lands on stigma is hydration. The pollen wall proteins are released from the surface. During the germination of pollen its entire content moves into the pollen tube. The growth is restricted to the tip of the tube and all the cytoplasmic contents move to the tip region. The remaining part of the pollen tube is occupied by a vacuole which is cut off from the tip by callose plug. The extreme tip of pollen tube appears hemispherical and transparent when viewed through the microscope. This is called **cap block**. As soon as the cap block disappear the growth of the pollen tube stops.

Pollen tube in the style

After the germination the pollen tube enters into the style from the stigma. The growth of the pollen tube in the style depends on the type of style.

Types of style

There are three types of style a) Hollow or open style b)solid style or closed style c) semi-solid or half closed style.

Hollow style (Open style): It is common among monocots. A hollow canal running from the stigma to the base of the style is present. The canal is lined by a single layer of glandular canal cells (Transmitting tissue).They secrete mucilaginous substances. The pollen tube grows on the surface of the cells lining the stylar canal. The canal is filled with secretions which serve as nutrition for growing pollen tubes and also controlling incompatibility reaction between the style and pollen tube. The secretions contain carbohydrates, lipids and some enzymes

TN_GOVT_BOTANY_XII_PAGES_001-032 CH 01.indd 20 23-12-2022 10:56:52