
Pollen calendar shows the production of pollen by plants during different seasons**.** This benefits the allergic persons. Pollen grains cause allergic reactions like asthma, bronchitis, hay fever, allergic rhinitis etc.,

Parthenium hysterophorus L. (Family-Asteraceae) is commonly called Carrot grass is a native of tropical America and was introduced into India as a contaminant along with cereal wheat. The pollen of this plant cause Allergy.



- The receptacle becomes fleshy and edible around the fruit enclosing the seeds as in *Pyrus malus* (apple)
- The calyx may persist and enlarge (*Solanum melongena)* or may cover the fruit *(Physalis minima*)



Physalis - Persistent calyx

- The flower stalk or axis below the gynoecium enlarges into a juicy pear shaped body which is edible (*Anacardium occidentale).* The Perianth becomes fleshy as in Jack fruit.



Anacardium - pedicel (edible)

- The cells present at the tip of the outer integument around the micropyle develop into a fleshy structure called **caruncle**. (*Ricinus communis).*



Ricinus - Caruncle

- The funiculus develops into a fleshy structure which is often very colourful and called **aril**. (*Myristica and Pithecellobium*)



Aril



Myristica *Pithecellobium*

- The nucellar tissue is either absorbed completely by the developing embryo sac and embryo or small portion may remain as storage tissue. Thus the remnant of nucellar tissue in the seed is called **perisperm**. Example: Black pepper and beet root

TN_GOVT_BOTANY_XII_PAGES_001-032 CH 01.indd 32 23-12-2022 10:57:01