- When the F_1offsprings are crossed with the dominant parents all the F_2 develop dominant character and no recessive individuals are obtained in the progeny.
- If the F_1 hybrid is crossed with the recessive parent individuals of both the phenotypes appear in equal proportion and this cross is specified as test cross.
- The recessive back cross helps to identify the heterozygosity of the hybrid.

2.3.4 Dihybrid cross

It is a genetic cross which involves individuals differing in two characters. Dihybrid inheritance is the inheritance of two separate genes each with two alleles.

Law of Independent Assortment – When two pairs of traits are combined in a hybrid, segregation of one pair of characters is independent to the other pair of characters. Genes that are located in different chromosomes assort independently during meiosis. Many possible combinations of factors can occur in the gametes.




Figure 2.9: Dihybrid cross – Segregation of gametes

Independent assortment leads to genetic diversity. If an individual produces genetically dissimilar gametes it is the consequence of independent assortment. Through independent assortment, the maternal and paternal members of all pairs were distributed to gametes, so all possible chromosomal combinations were produced leading to genetic variation. In sexually reproducing plants / organisms, due to independent assortment, genetic variation takes place which is important in the process of evolution. The Law of Segregation is concerned with alleles of one gene but the Law of Independent Assortment deals with the relationship between genes.

The crossing of two plants differing in two pairs of contrasting traits is called dihybrid cross. In dihybrid cross, two characters (colour and shape) are considered at a time. Mendel considered the seed shape (round and wrinkled) and cotyledon colour (yellow & green) as the two characters. In seed shape round (R) is dominant over wrinkled (r) ; in cotyledon colour yellow (Y) is dominant over green (y). Hence the pure breeding round yellow parent is represented by the genotype RRYY and the pure breeding green wrinkled parent is represented by the genotype rryy. During gamete formation the paired genes of a character assort out independently of the other pair. During the F_1 x F_1 fertilization each zygote with an equal probability receives one of the four combinations from each parent. The resultant gametes thus will be genetically different and they are of the following four types:

1) Yellow round (YR) - 9/16
2) Yellow wrinkled (Yr) - 3/16
3) Green round (yR) - 3/16
4) Green wrinkled (yr) - 1/16

These four types of gametes of F_1 dihybrids unite randomly in the process of fertilization and produce sixteen types of individuals in F_2 in the ratio of 9:3:3:1 as shown in the figure. Mendel's 9:3:3:1 dihybrid ratio is an ideal ratio based on the probability including segregation, independent assortment and random

TN_GOVT_BOTANY_XII_PAGES_033-051 CH 02.indd 40 23-12-2022 10:57:36