substitutions and base pair insertions or deletions. Base substitutions are mutations in which there is a change in the DNA such that one base pair is replaced by another. It can be divided into two subtypes: transitions and transversions. Addition or deletion mutations are actually additions or deletions of nucleotide pairs and also called base pair addition or deletions. Collectively, they are termed **indel mutations** (for insertion-deletion).

Substitution mutations or indel mutations affect translation. Based on these different types of mutations are given below.

The mutation that changes one codon for an amino acid into another codon for that same amino acid are called **Synonymous or silent mutations.** The mutation where the codon for one amino acid is changed into a codon for another amino acid is called **Missense or non-synonymous mutations**. The mutations where codon for one amino acid is changed into a termination or stop codon is called **Nonsense mutation**. Mutations that result in the addition or deletion of a single base pair of DNA that changes the reading frame for the translation process as a result of which there is complete loss of normal protein structure and function are called **Frameshift mutations** (Figure: 3.9).

3.5.2 Mutagenic agents

The factors which cause genetic mutation are called **mutagenic agents or mutagens**. Mutagens are of two types, physical mutagen and chemical mutagen. **Muller** (1927) was the first to find out physical mutagen in *Drosophila*.

Physical mutagens:

Scientists are using temperature and radiations such as X rays, gamma rays, alfa rays, beta rays, neutron, cosmic rays, radioactive isotopes, ultraviolet rays as physical mutagen to produce mutation in various plants and animals.

Temperature: Increase in temperature increases the rate of mutation. While rise in temperature, breaks the hydrogen bonds between two DNA nucleotides which affects the process of replication and transcription.

Radiation: The electromagnetic spectrum contains shorter and longer wave length rays than the visible spectrum. These are classified into ionizing and non-ionizing radiation. Ionizing radiation are short wave length and carry enough higher energy to ionize electrons from atom. X rays, gamma rays, alfa rays, beta rays and cosmic rays which breaks the chromosomes (chromosomal mutation) and chromatids in irradiated cells. Non-ionizing radiation, UV rays have longer wavelengths and carry lower energy, so they have lower penetrating power than the ionizing radiations. It is used to treat unicellular microorganisms, spores, pollen grains which possess nuclei located near surface membrane.

Sharbati Sonora

Sharbati Sonora is a mutant variety of wheat, which is developed from Mexican variety (Sonora 64) by irradiating of gamma rays. It is the work of Dr. **M.S.Swaminathan** who is known as **'Father of Indian green revolution'** and his team.

Castor Aruna

Castor Aruna is mutant variety of castor which is developed by treatment of seeds with thermal neutrons in order to induce very early maturity (120 days instead of 270 days as original variety).

Chemical mutagens:

Chemicals which induce mutation are called chemical mutagens. Some chemical mutagens are mustard gas, nitrous acid, ethyl and methyl methane sulphonate (EMS and MMS), ethyl urethane, magnous salt, formaldehyde, eosin and enthrosine. Example: Nitrous oxide alters the nitrogen bases of DNA and disturb the replication and transcription

TN_GOVT_BOTANY_XII_PAGES_052-081 CH 03.indd 62 28-12-2022 15:41:18