


Figure 3.10 Types of Ploidy

respectively. All possible tetrasomics are available in Wheat.

(c) Pentasomy

Addition of three individual chromosome from different chromosomal pairs to normal diploid set are called pentasomy (2n+3).

2. Hypoploidy

Loss of one or more chromosome from the diploid set in the cell is called **hypoploidy**. It can be divided into two types. They are

(a) Monosomy

Loss of a single chromosome from the diploid set are called **monosomy**(2n-1). However loss of two individual or three individual chromosomes are called **double monosomy** (2n-1-1) and **triple monosomy** (2n-1-1-1) respectively. Double monosomics are observed in maize.

(b) Nullisomy

Loss of a pair of homologous chromosomes or two pairs of homologous chromosomes from the diploid set are called **Nullisomy** (2n-2) and **double Nullisomy** (2n-2-2) respectively. Selfing of monosomic plants produce nullisomics. They are usually lethal.




Figure 3.11 Types of aneuploidy

(ii) Euploidy

Euploidy is a condition where the organisms possess one or more basic sets of chromosomes. Euploidy is classified as monoploidy, diploidy and polyploidy. The condition where an organism or somatic cell has two sets of chromosomes are called diploid (2n). Half the number of somatic chromosomes is referred as gametic chromosome number called haploid(n). It should be noted that haploidy (n) is different from a monoploidy (x). For example, the common wheat plant

TN_GOVT_BOTANY_XII_PAGES_052-081 CH 03.indd 64 28-12-2022 15:41:18