



**Figure 4.13:** Liposome mediated method of Gene Transfer





**Figure 4.14:** Gene gun method of Gene Transfer

or tissues are cultured on selected medium to regenerate plants from the transformed cells.(Figure 4.14)

### 4.5.2 Indirect or Vector-Mediated Gene Transfer

Gene transfer is mediated with the help of a plasmid vector is known as indirect or vector mediated gene transfer. Among the various vectors used for plant transformation, the Ti-plasmid from *Agrobacterium tumefaciens* has been used extensively. This bacterium has a large size plasmid, known as Ti plasmid (Tumor inducing) and a portion of it referred as T-DNA (transfer DNA) is transferred to plant genome in the infected cells and cause plant tumors (crown gall). Since this bacterium has the natural ability to transfer T-DNA region of its plasmid into plant genome, upon infection of cells at the wound site, it is also known as the natural genetic engineer of plants.

The foreign gene (e.g. Bt gene for insect resistance) and plant selection marker gene, usually an antibiotic gene like *npt II* which confers resistance to antibiotic kanamycin are cloned in the T DNA region of Ti-plasmid in place of unwanted DNA sequences.(Figure 4.15)

## 4.6 Screening for Recombinants

After the introduction of *r*-DNA into a suitable host cell, it is essential to identify those cells which have received the r-DNA molecule. This process is called screening. The vector or foreign DNA present in recombinant cells expresses the characters, while the non-recombinants do not express the characters or traits. For this some of





**Figure 4.15:** Agrobacterium mediated gene transfer in plants



TN_GOVT_BOTANY_XII_PAGES_082-109 CH 04.indd 91 23-12-2022 10:58:54