radiant energy, CO_2, O_2 and space, etc.

6. **Reaction** - The species occupying a habitat gradually modify the environmental condition, where the existing species community is displaced or replaced by another. This is called reaction. The community which is replaced by another community is called **seral community**.

7. Stabilization (Climax stage) - The final establishment of plant community is called stabilization. This establishment of a plant community which maintains itself in equilibrium with climax of the area and not replaced by others is known as **climax community** and the stage is climax stage.

7.3.5 Classification of plant succession

Detailed study of Hydrosere and Lithosere are discussed below:




Figure 7.19: Classification of plant succession

Hydrosere

The succession in a freshwater ecosystem is also referred to as hydrosere. Succession in a pond, begins with colonization of the pioneers like phytoplankton and finally ends with the formation of climax community like forest stage. It includes the following stages Fig 7.20.

1. Phytoplankton stage - It is the first stage of succession consisting of the pioneer community like blue green algae, green algae, diatoms, bacteria, etc., The colonization of these organisms enrich the amount of organic matter and nutrients of pond due to their life activities and death. This favors the development of the next seral stages.

2. Submerged plant stage - As the result of death and decomposition of planktons, silt brought from land by rain water, lead to a loose mud formation at the bottom of the pond. Hence, the rooted submerged hydrophytes begin to appear on the new substratum. Example: *Chara, Vallisneria* and *Hydrilla* etc. The death and decay of these plants will build up the substratum of pond to become shallow. Therefore, this habitat now replaces another group of plants which are of floating type.

3. Submerged free floating stage - During this stage, the depth of the pond will become almost 2-5 feet. Hence, the rooted hydrophytic plants and with floating large leaves start colonising the pond. Example: Rooted floating plants like *Nelumbo, Nymphaea* and *Trapa*. Some free floating species like *Azolla, Lemna, Wolffia* and *Pistia* are also present in this stage. By death and decomposition of these plants, further the pond becomes more shallow. Due to this reason, floating plant species is gradually replaced by another species which makes new seral stage.

4. Reed-swamp stage - It is also called an amphibious stage. During this stage, rooted floating plants are replaced by plants which can live successfully in aquatic as well as aerial environment. Example: *Typha, Phragmites, Sagittaria* and *Scirpus* etc. At the end of this stage, water level is very much reduced, making it unsuitable for the continuous growth of amphibious plants.

5. Marsh meadow stage - When the pond becomes swallowed due to decreasing water level, species of Cyperaceae and Poaceae such as *Carex, Juncus, Cyperus* and *Eleocharis* colonise

TN_GOVT_BOTANY_XII_PAGES_149-177 CH 07.indd 165 23-12-2022 11:03:04