Titration –I

(Link $KMnO_4$)Vs (Standard $FeSO_4$)

s.no	Volume of standard $FeSO_4$ (ml)	Burette readings		Concordant value (Volume of $KMnO_4$) (ml)
		Initial (ml)	Final (ml)	
1	20			
2	20			
3	20			

Calculation :

Volume of $KMnO_4$ (link) solution V_1 = ml

Normality $KMnO_4$ (link) solution N_1 = ?N

Volume of standard $FeSO_4$ solution V_2 = 20 ml

Normality of standard $FeSO_4$ solution N_2 = 0.1024 N

According to normality equation:

According to normality equation: $V_1 \times N_1 = V_2 \times N_2$

$$N_1 = \frac{V_2 \times N_2}{V_1}$$

Normality of $KMnO_4$ (link) solution (N_1) = _____X_____ N

Titration–II

(Unknown FAS) Vs (Link $KMnO_4$)

Burette is washed with water, rinsed with $KMnO_4$ solution and filled with same $KMnO_4$ solution up to the zero mark. Exactly 20 ml of unknown FAS solution is pipetted out into the clean, washed conical flask. To this FAS solution approximately 20ml of 2N sulphuric acid is added. This mixture is titrated against $KMnO_4$ Link solution from the burette. $KMnO_4$ is added drop wise till the appearance of permanent pale pink colour. Burette reading is noted and the same procedure is repeated to get concordant values.

organic analysis.indd 269 22-12-2022 19:48:37