Titration –II

(Link $KMnO_4$)Vs (Unknown FAS)

s.no	Volume of Unknown FAS (ml)	Burette readings		Concordant value (Volume of $KMnO_4$) (ml)
		Initial (ml)	Final (ml)	
1	20			
2	20			
3	20			

Calculation :

Volume of Unknown FAS solution V_1 = 20ml

Normality of Unknown FAS solution N_1 = ? N

Volume of $KMnO_4$ (link) solution V_2 = ml

Normality $KMnO_4$ (link) solution N_2 = N

According to normality equation: $V_1 \times N_1 = V_2 \times N_2$

$$N_1 = \frac{V_2 \times N_2}{V_1}$$

$$N_1 = \frac{Y}{\quad\quad} N$$

The normality of unknown FAS solution $= \frac{Y}{\quad\quad} N$

Weight calculation:

The amount of FAS dissolved in 1 lit of the solution = (Normality) x (equivalent weight)

The amount of FAS dissolved in 1500 ml of the solution $= \frac{\text{Normality} \times \text{equivalentweight} \times 1500}{1000}$

$$= \frac{Y \times 392 \times 1500}{1000}$$

$$= \quad g$$

Report :

The amount of FAS dissolved in 1500 ml of the solution = g

organic analysis.indd 270 22-12-2022 19:48:37