Titration –II

(Link $KMnO_4$)Vs (Unknown oxalic acid)

s.no	Volume of Unknown oxalic acid (ml)	Burette readings		Concordant value (Volume of $KMnO_4$) (ml)
		Initial (ml)	Final (ml)	
1	20			
2	20			
3	20			

Calculation :

Volume of Unknown oxalic acid solution V_1 = 20 ml

Normality of Unknown oxalic acid solution N_1 = ? N

Volume of $KMnO_4$ (link) solution V_2 = ml

Normality $KMnO_4$ (link) solution N_2 = N

According to normality equation:

$$V_1 \times N_1 = V_2 \times N_2$$

$$N_1 = \frac{V_2 \times N_2}{V_1}$$

Normality of Unknown oxalic acid solution N_1 = $\frac{Y}{\quad\quad}$ N

Weight calculation:

The amount of oxalic acid dissolved in 1 lit of the solution =(Normality) x (equivalent weight)

The amount of oxalic acid dissolved in 500 ml of the solution $= \frac{Y \times 63 \times 500}{1000}$

$$= \frac{\quad \text{x } 63 \text{ x } 500}{1000}$$

$$= \quad \text{g}$$

Report :

The amount of oxalic acid dissolved in 500 ml of given the solution = g

organic analysis.indd 273 22-12-2022 19:48:38