Titration –I

(standard HCl)Vs **(link** NaOH)

s.no	Volume of NaOH(ml)	Burette readings		Concordant value (Volume of std HCl) (ml)
		Initial (ml)	Final (ml)	
1	20			
2	20			
3	20			

Calculation :

Volume of NaOH(link) solution $V_1 = 20$ ml

Normality NaOH(link) solution $N_1 = ?$ N

Volume of standard HCl solution $V_2 =$ ml

Normality of standard HCl solution $N_2 = 0.1010$ N

According to normality equation:

$$V_1 \times N_1 = V_2 \times N_2$$

$$N_1 = \frac{\times\ 0.1010}{20} =$$

Normality NaOH (link) solution $N_1 =$ _____X_____ N

Titration–II

(Unknown oxalic acid) Vs (Link NaOH)

Burette is washed with water, rinsed with oxalic acid solution and filled with same oxalic acid solution up to the zero mark. Exactly 20 ml of NaOH solution is pipetted out into the clean, washed conical flask. To This solution 2 to 3 drops of phenolphthalein indicator is added and titrated against oxalic acid solution from the burette. oxalic acid is added drop wise till the pink colour disappears completely. Burette reading is noted and the same procedure is repeated to get concordant values.

organic analysis.indd 278 22-12-2022 19:48:39