Titration –II

(Link NaOH)Vs (Unknown oxalic acid solution)

s.no	Volume of NaOH link (ml)	Burette readings		Concordant value (Volume of oxalic acid) (ml)
		Initial (ml)	Final (ml)	
1	20			
2	20			
3	20			

Calculation :

Volume of Unknown oxalic acid solution V_1 = ml

Normality of Unknown oxalic acid solution N_1 = ? N

Volume of NaOH solution V_2 = 20 ml

Normality NaOH solution N_2 = N

According to normality equation:

$$V_1 \times N_1 = V_2 \times N_2$$

$$N_1 = \frac{V_2 \times N_2}{V_1}$$

Normality of Unknown oxalic acid solution $N_1 = \frac{Y}{_____} N$

Weight calculation:

The amount of oxalic acid dissolved in 1 lit of the solution = (Normality) x (equivalent weight)

The amount of oxalic acid dissolved in 1250 ml of the solution $= \frac{\text{Normality x equivalentweight x 1250}}{1000}$

$$= \frac{Y \times 63 \times 1250}{1000}$$

$$= \frac{\text{x 63 x 1250}}{1000}$$

$$= \quad \text{g}$$

Report :

The amount of oxalic acid dissolved in 1250 ml of the solution = g

organic analysis.indd 279 22-12-2022 19:48:39