
When the bar magnet is cut along the axis into two pieces, new magnetic pole strength is $q'_m = \frac{q_m}{2}$ but magnetic length does not change. So, the magnetic moment is


$$p'_m = q'_m 2l$$

$$p'_m = \frac{q_m}{2} 2l = \frac{1}{2}(q_m 2l) = \frac{1}{2} p_m$$

In vector notation, $\vec{p}'_m = \frac{1}{2}\vec{p}_m$

(b) a bar magnet cut into two pieces perpendicular to the axis:






When the bar magnet is cut perpendicular to the axis into two pieces, magnetic pole strength will not change but magnetic length will be halved. So the magnetic moment is

$$p'_m = q_m \times \frac{1}{2}(2l) = \frac{1}{2}(q_m \cdot 2l) = \frac{1}{2} p_m$$

In vector notation, $\vec{p}'_m = \frac{1}{2}\vec{p}_m$

EXAMPLE 3.3

Compute the magnetic length of a uniform bar magnet if the geometrical length of the magnet is 12 cm. Mark the positions of magnetic pole points.




Solution

Geometrical length of the bar magnet is 12 cm

$$\text{Magnetic length} = \frac{5}{6} \times (\text{geometrical length})$$

$$= \frac{5}{6} \times 12 = 10\,\text{cm}$$

In this figure, the dot implies the pole points.



Note

(i) Pole strength is a scalar quantity with dimension $[M^0LT^0A]$. Its SI unit is NT^{-1} (newton per tesla) or A m (ampere-metre).

(ii) Like positive and negative charges in electrostatics, north pole of a magnet experiences a force in the direction of magnetic field while south pole of a magnet experiences force opposite to the magnetic field.

(iii) Pole strength depends on the nature of materials of the magnet, area of cross-section and the state of magnetization.

(iv) If a magnet is cut into two equal halves along the length then pole strength is reduced to half.

(v) If a magnet is cut into two equal halves perpendicular to the length, then pole strength remains same.

(vi) If a magnet is cut into two pieces, we will not get separate north and south poles. Instead, we get two magnets. In other words, isolated monopole does not exist in nature.

12th Physics_Vol-1_Unit 3_English Medium.indd 132 23-12-2022 15:01:31