elemental copper is more stable than Cu^{2+}. There are two deviations., In the general trend, Fig shows that $\left(E^{0}_{M^{2+}/M}\right)$ value for manganese and zinc are more negative than the regular trend. It is due to extra stability which arises due to the half filled d^5 configuration in Mn^{2+} and completely filled d^{10} configuration in Zn^{2+}.


Transition metals in their high oxidation states tend to be oxidizing . For example, Fe^{3+} is moderately a strong oxidant, and it oxidises copper to Cu^{2+} ions. The feasibility of the reaction is predicted from the following standard electrode potential values.

$Fe^{3+}(aq) + e^{-} \rightleftharpoons Fe^{2+} \quad E^{0} = 0.77V$

$Cu^{2+}(aq) + 2e^{-} \rightleftharpoons Cu(s) \quad E^{0} = +0.34\ V$

The standard electrode potential for the M^{3+}/M^{2+} half-cell gives the relative stability between M^{3+} and M^{2+}. The reduction potential values are tabulated as below.

Reaction	Standard reduction potential (V)
$Ti^{3+} + e^{-} \longrightarrow Ti^{2+}$	–0.37
$V^{3+} + e^{-} \longrightarrow V^{2+}$	–0.26
$Cr^{3+} + e^{-} \longrightarrow Cr^{2+}$	–0.41
$Mn^{3+} + e^{-} \longrightarrow Mn^{2+}$	+1.51
$Fe^{3+} + 2e^{-} \longrightarrow Fe^{2+}$	+0.77
$Co^{3+} + 2e^{-} \longrightarrow Co^{2+}$	+1.81



The negative values for titanium, vanadium and chromium indicate that the higher oxidation state is preferred. If we want to reduce such a stable Cr^{3+} ion, strong reducing agent which has high negative value for reduction potential like metallic zinc $\left(E^{0} = -0.76\ V\right)$ is required.

The high reduction potential of Mn^{3+}/Mn^{2+} indicates Mn^{2+} is more stable than Mn^{3+}. For Fe^{3+}/Fe^{2+} the reduction potential is 0.77V, and this low value indicates that both Fe^{3+} and Fe^{2+} can exist under normal conditions. The drop from Mn to Fe is due to the electronic structure of the ions concerned.Mn^{3+} has a $3d^4$ configuration while that of Mn^{2+} is $3d^5$. The extra stability associated with a half filled d sub shell makes the reduction of Mn^{3+} very feasible ($E^{0} = +1.51V$).

4.3.6 Magnetic properties

Most of the compounds of transition elements are paramagnetic. Magnetic properties are related to the electronic configuration of atoms. We have already learnt in XI STD that

XII U4-D-Block-Jerald Folder.indd 108 3/29/2019 6:46:33 PM