3. Among the transition metals of 3d series, the one that has highest negative $\left(M^{2+}/M\right)$ standard electrode potential is

   a) Ti b) Cu c) Mn d) Zn

4. Which one of the following ions has the same number of unpaired electrons as present in V^{3+}?

   a) Ti^{3+} b) Fe^{3+}

   c) Ni^{2+} d) Cr^{3+}

5. The magnetic moment of Mn^{2+} ion is

   a) 5.92BM b) 2.80BM

   c) 8.95BM d) 3.90BM

6. Which of the following compounds is colourless?

   a) Fe^{3+} b) Ti^{4+}

   c) Co^{2+} d) Ni^{2+}

7. the catalytic behaviour of transition metals and their compounds is ascribed mainly due to

   a) their magnetic behaviour

   b) their unfilled d orbitals

   c) their ability to adopt variable oxidation states

   d) their chemical reactivity

8. The correct order of increasing oxidizing power in the series

   a) $VO_2^+ < Cr_2O_7^{2-} < MnO_4^-$ b) $Cr_2O_7^{2-} < VO_2^+ < MnO_4^-$

   c) $Cr_2O_7^{2-} < MnO_4^- < VO_2^+$ d) $MnO_4^- < Cr_2O_7^{2-} < VO_2^+$

9. The alloy of copper that contain Zinc is

   a) Monel metal b) Bronze

   c) bell metal d) brass

10. Which of the following does not give oxygen on heating?

    a) $K_2Cr_2O_7$ b) $(NH_4)_2Cr_2O_7$

    c) $KClO_3$ d) $Zn(ClO_3)_2$

11. In acid medium, potassium permanganate oxidizes oxalic acid to

    a) oxalate b) Carbon dioxide

    c) acetate d) acetic acid

XII U4-D-Block-Jerald Folder.indd 126 3/29/2019 6:46:53 PM