13. A solid compound XY has NaCl structure. if the radius of the cation is 100pm , the radius of the anion will be

a) $\left(\frac{100}{0.414}\right)$ b) $\left(\frac{0.732}{100}\right)$ c) 100×0.414 d) $\left(\frac{0.414}{100}\right)$

14. The vacant space in bcc lattice unit cell is

a) 48% b) 23% c) 32% d) 26%

15. The radius of an atom is 300pm, if it crystallizes in a face centered cubic lattice, the length oif the edge of the unit cell is

a) 488.5pm b) 848.5pm c) 884.5pm d) 484.5pm

16. The fraction of total volume occupied by the atoms in a simple cubic is

a) $\left(\frac{\pi}{4\sqrt{2}}\right)$ b) $\left(\frac{\pi}{6}\right)$ c) $\left(\frac{\pi}{4}\right)$ d) $\left(\frac{\pi}{3\sqrt{2}}\right)$

17. The yellow colour in NaCl crystal is due to

a) excitation of electrons in F centers

b) reflection of light from Cl^- ion on the surface

c) refraction of light from Na^+ ion

d) all of the above

18. if 'a' stands for the edge length of the cubic system ; sc , bcc, and fcc. Then the ratio of radii of spheres in these systems will be respectively.

a) $\left(\frac{1}{2}a : \frac{\sqrt{3}}{2}a : \frac{\sqrt{2}}{2}a\right)$ b) $\left(\sqrt{1}a : \sqrt{3}a : \sqrt{2}a\right)$

c) $\left(\frac{1}{2}a : \frac{\sqrt{3}}{4}a : \frac{1}{2\sqrt{2}}a\right)$ d) $\left(\frac{1}{2}a : \sqrt{3}a : \frac{1}{\sqrt{2}}a\right)$

19. if 'a' is the length of the side of the cube, the distance between the body centered atom and one corner atom in the cube will be

a) $\left(\frac{2}{\sqrt{3}}\right)a$ b) $\left(\frac{4}{\sqrt{3}}\right)a$

c) $\left(\frac{\sqrt{3}}{4}\right)a$ d) $\left(\frac{\sqrt{3}}{2}\right)a$

20. Potassium has a bcc structure with nearest neighbor distance 4.52 A^0 . its atomic weight is 39. its density will be

a) 915 kg m^{-3} b) 2142 kg m^{-3} c) 452 kg m^{-3} d) 390 kg m^{-3}

21. Schottky defect in a crystal is observed when

a) unequal number of anions and anions are missing from the lattice

XII U6 Solid State - Jerald.indd 199 3/29/2019 6:30:38 PM