Titration –II

(Link $KMnO_4$)Vs (Unknown oxalic acid)

s.no	Volume of Unknown oxalic acid (ml)	Burette readings		Concordant value (Volume of $KMnO_4$) (ml)	indicator
		Initial (ml)	Final (ml)		
1	20	0			Self (1)
2	20	0			
3	20	0			

Calculation :

Volume of Unknown oxalic acid solution N_1 = 20 ml

Normality of Unknown oxalic acid solution N_1 = ? N

Volume of 2 (link) solution V_2 = ml

Normality 2 (link) solution N_2 = N

According to normality equation:

$$V_1 \times N_1 = V_2 \times N_2$$

$$N_1 = \frac{V_2 \times N_2}{V_1}$$

$$N_1 = \frac{20 \times 0.1024}{} =$$

Normality of Unknown oxalic acid solution N_1 = _____ N

Weight calculation:

The amount of oxalic acid dissolved in 1 lit of the solution = (Normality) x (equivalent weight)

The amount of oxalic acid dissolved in 500 ml of the solution = $\frac{\text{x 63 x 500}}{1000}$

= $\frac{}{1000}$

= g

Report :

The normality of $KMnO_4$ link solution = N

The normality of unknown oxalic acid solution = N

The amount of oxalic acid dissolved in 500 ml of given the solution = g

organic analysis.indd 273 3/29/2019 6:48:27 PM