Titration –I

(standard HCl)Vs **(link** NaOH)

s.no	Volume of NaOH(ml)	Burette readings		Concordant value (Volume of std HCl) (ml)	Indicator
		Initial (ml)	Final (ml)		
1	20	0			phenolphthalein
2	20	0			
3	20	0			

Calculation :

Volume of NaOH(link) solution V_1 = 20 ml

Normality NaOH(link) solution V_2 = ? N

Volume of standard HCl solution V_2 = ml

Normality of standard HCl solution N_2 = 0.1010 N

According to normality equation:

$$V_1 \times N_1 = V_2 \times N_2$$

$$N_1 = \frac{\times\ 0.1010}{20} =$$

$$N_1 = \frac{\times\ 0.1010}{20} =$$

Normality NaOH (link) solution _____ N

Titration–II

(Unknown oxalic acid) Vs (Link NaOH)

Burette is washed with water, rinsed with oxalic acid solution and filled with same oxalic acid solution up to the zero mark. Exactly 20 ml of NaOH solution is pipetted out into the clean, washed conical flask. To This solution 2 to 3 drops of phenolphthalein indicator is added and titrated against oxalic acid solution from the burette. oxalic acid is added drop wise till the pink colour disappears completely. Burette reading is noted and the same procedure is repeated to get concordant values.

organic analysis.indd 278 3/29/2019 6:48:28 PM