Titration –II

(Link NaOH)Vs (Unknown oxalic acid solution)

s.no	Volume of NaOH link (ml)	Burette readings		Concordant value (Volume of oxalic acid) (ml)	Indicator
		Initial (ml)	Final (ml)		
1	20	0			Phenolphthalein
2	20	0			
3	20	0			

Calculation :

Volume of Unknown oxalic acid solution N_1 = ml

Normality of Unknown oxalic acid solution N_1 = ? N

Volume of NaOH solution N_2 = 20 ml

Normality NaOH solution N_2 = N

According to normality equation:

V_1

$$N_1 = \frac{V_2 \times N_2}{V_1}$$

$$\frac{\text{Normality x equivalentweight x 1250}}{1000}$$

Weight calculation:

The amount of oxalic acid dissolved in 1 lit of the solution = (Normality) x (equivalent weight)

The amount of oxalic acid dissolved in 1250 ml of the solution $= \frac{\text{Normality x equivalentweight x 1250}}{1000}$

$= \frac{\text{x 63 x 1250}}{1000}$

= g

Report :

1. The normality of NaOH Link solution = N
2. The normality of unknown oxalic acid solution = N
3. The amount of oxalic acid dissolved in 1250 ml of the solution = g

organic analysis.indd 279 3/29/2019 6:48:28 PM