Table 8.2 Differences between active and passive immunity

Sl.No	Active Immunity	Passive Immunity
1	Active immunity is produced actively by host's immune system.	Passive immunity is received passively and there is no active host participation.
2	It is produced due to contact with pathogen or by its antigen.	It is produced due to antibodies obtained from outside.
3	It is durable and effective in protection.	It is transient and less effective.
4	Immunological memory is present.	No memory.
5	Booster effect on subsequent dose is possible.	Subsequent dose is less effective.
6	Immunity is effective only after a short period.	Immunity develops immediately.

immune system for **the first time**. During this, the immune system has to learn to recognize the antigen, produce antibody against it and eventually produce memory lymphocytes. The primary immune response is slow and short-lived.

Secondary immune response

The secondary immune response occurs when a person is exposed to the same antigen again. During this time, immunological memory has been established and the immune system can start producing antibodies **immediately**. Within hours after recognition of the antigen, a new army of plasma cells are generated. Within 2 to 3 days, the antibody concentration in the blood **rises steeply** to reach much higher level than primary response. This is also called as "**booster response**".

8.5 Lymphoid organs

Immune system of an organism consists of several structurally and functionally different organs and tissues that are widely dispersed in the body. The organs involved in the origin, maturation and proliferation of lymphocytes are called **lymphoid organs (Fig. 8.3).**

Table 8.3 Differences between primary and secondary immune responses

Sl.No	Primary Immune Response	Secondary Immune Response
1	It occurs as a result of primary contact with an antigen.	It occurs as a result of second and subsequent contacts with the same antigen.
2	Antibody level reaches peak in 7 to 10 days.	Antibody level reaches peak in 3 to 5 days.
3	Prolonged period is required to establish immunity.	It establishes immunity in a short time.
4	There is rapid decline in antibody level.	Antibody level remains high for longer period.
5	It appears mainly in the lymph nodes and spleen.	It appears mainly in the bone marrow, followed by the spleen and lymph nodes.

XII Std Zoology Chapter 8 EM.indd 132 23/12/2022 16:43:14