synthesis the temperature of the mixture is increased to 75^{0}C for a sufficient period of time to allow Taq DNA polymerase to extend each primer by copying the single stranded template. At the end of incubation both single template strands will be made partially double stranded. The new strand of each double stranded DNA extends to a variable distance downstream. These steps are repeated again and again to generate multiple forms of the desired DNA. This process is also called DNA amplification.

The PCR technique can also be used for amplifications of RNA in which case it is referred to as reverse transcription PCR (RT-PCR). In this process the RNA molecules (mRNA) must be converted to complementary DNA by the enzyme reverse transcriptase. The cDNA then serves as the template for PCR.

PCR In Clinical Diagnosis

The specificity and sensitivity of PCR is useful for the diagnosis of inherited disorders (genetic diseases), viral diseases, bacterial diseases, etc., The diagnosis and treatment of a particular disease often requires identifying a particular pathogen. Traditional methods of identification involve culturing these organisms from clinical specimens and performing metabolic and other tests to identify them. The concept behind PCR based diagnosis of infectious diseases is simple – if the pathogen is present in a clinical specimen its DNA will be present. Its DNA has unique sequences that can be detected by PCR, often using the clinical specimen (for example, blood, stool, spinal fluid, or sputum) in the PCR mixture. PCR is also employed in the prenatal diagnosis of inherited diseases by using chorionic villi samples or cells from amniocentesis. Diseases like sickle cell anemia, β-thalassemia and phenylketonuria can be detected by PCR in these samples. cDNA from PCR is a valuable tool for diagnosis and monitoring retroviral infections e.g., Corona Virus (SARS - CoV-2).

Several virally induced cancers, like cervical cancer caused by Papilloma virus can be detected by PCR. Sex of human beings and live stocks, embryos fertilized *invitro* can be determined by PCR by using primers and DNA probes specific for sex chromosomes. PCR technique is also used to detect sex-linked disorders in fertilized embryos.

Applications of PCR

The differences in the genomes of two different organisms can be studied by PCR. PCR is very important in the study of evolutions, more specifically phylogenetics.




Fig. 10.6 Steps involved in PCR

XII Std Zoology Chapter 10 EM.indd 165 23/12/2022 16:50:12