Table 7.5 Differences between active and passive immunity

Sl.No	Active Immunity	Passive Immunity
1	Active immunity is produced actively by host's immune system.	Passive immunity is received passively and there is no active host participation.
2	It is produced due to contact with pathogen or by its antigen.	It is produced due to antibodies obtained from outside.
3	It is durable and effective in protection.	It is transient and less effective.
4	Immunological memory is present.	No memory.
5	Booster effect on subsequent dose is possible.	Subsequent dose is less effective.
6	Immunity is effective only after a short period.	Immunity develops immediately.

The immunological resistance developed by the organisms through the production of antibodies in their body is called active immunity. Active immunity is acquired through the use of a person's immune responses, which lead to the development of memory cells. Active immunity results from an infection or an immunization.

Passive immunity does not require the body to produce antibodies to antigens. The antibodies are introduced from outside into the organism. Thus, passive immunity is acquired without the activation of a person's immune response, and therefore there is no memory.

7.3.3 Immune responses

The immune responses may be **primary** or **secondary (Table 7.6)**.



The process of production of blood cells in the bone marrow is called **haematopoiesis**.

Primary immune response

The primary immune response occurs when a pathogen comes in contact with the immune system for **the first time**. During this, the immune system has to learn to

Table 7.6 Differences between primary and secondary immune responses

Sl.No	Primary Immune Response	Secondary Immune Response
1	It occurs as a result of primary contact with an antigen.	It occurs as a result of second and subsequent contacts with the same antigen.
2	Antibody level reaches peak in 7 to 10 days.	Antibody level reaches peak in 3 to 5 days.
3	Prolonged period is required to establish immunity.	It establishes immunity in a short time.
4	There is rapid decline in antibody level.	Antibody level remains high for longer period.
5	It appears mainly in the lymph nodes and spleen.	It appears mainly in the bone marrow, followed by the spleen and lymph nodes.

XII Std Biology-Zoology Chapter-7 EM.indd 118 23/12/2022 12:39:56