nm wavelength of light. PAR values range from 0 to 3000 millimoles /square meter / second. At night PAR is zero and during midday in the summer, PAR often reaches 2000 – 3000 millimoles /square meter/second.



Types of Carbon

Green carbon – carbon stored in the biosphere (by the process of photosynthesis).

Grey carbon – carbon stored in fossil fuel (coal, oil and biogas deposits in the lithosphere).

Blue carbon – carbon stored in the atmosphere and oceans.

Brown carbon – carbon stored in industrialized forests (wood used in making commercial articles)

Black carbon – carbon emitted from gas, diesel engine and coal fired power plants.

7.2.2 Productivity of an ecosystem

The rate of biomass production per unit area in a unit time is called productivity. It can be expressed in terms of gm /m^2/year or Kcal/m^2/ year. It is classified as given bellow.

1. Primary productivity
2. Secondary productivity
3. Community productivity

1. Primary productivity:

The chemical energy or organic matter generated by autotrophs during the process of photosynthesis and chemosynthesis is called **primary productivity**. It is the source of energy for all organisms, from bacteria to human.

a. Gross Primary Productivity (GPP)

The total amount of food energy or organic matter or biomass produced in an ecosystem by autotrophs through the process of photosynthesis is called **gross primary productivity**

b. Net Primary Productivity (NPP)

The proportion of energy which remains after respiration loss in the plant is called **net primary productivity**. It is also called as apparent photosynthesis. Thus the difference between GPP and respiration is known as NPP.

$$NPP = GPP - Respiration$$

NPP of whole biosphere is estimated to be about 170 billion tons (dry weight) per year. Out of which NPP of oceanic producers is only 55 billion tons per year.

2. Secondary productivity

The amount of energy stored in the tissues of heterotrophs or consumers is called **secondary productivity**.

a. Gross secondary productivity

It is equivalent to the total amount of plant material ingested by the herbivores minus the materials lost as faeces.

b. Net secondary productivity

Storage of energy or biomass by consumers per unit area per unit time, after respiratory loss is called **net secondary productivity**.

3. Community productivity

The rate of net synthesis of organic matter (biomass) by a group of plants per unit area per unit time is known as **community productivity**.

Factors affecting primary productivity

Primary productivity depends upon the plant species of an area, their photosynthetic capacity, availability of nutrients, solar radiation, precipitation, soil type, topographic factors (altitude, latitude, direction), and other environmental factors. It varies in different types of ecosystems.

7.2.3 Concept of trophic level in an ecosystem

(Greek word ' trophic' = to food or feeding)

A trophic level refers to the position of an organism in the food chain. The number of

TN_GOVT_BOTANY_XII_PAGES_149-177 CH 07.indd 151 23-12-2022 11:01:28