35. The technique of obtaining large number of plantlets by tissue culture method is called
a) Plantlet culture (AIPMT 2005)
b) Organ culture
c) Micropropagation
d) Macropropagation

36. Coconut milk is used in tissue culture in which present (AIPMT 2000)
a) cytokinin b) auxin
c) gibberellins d) ethylene.

37. Haploid plants can be obtained by culturing. (AIPMT 1994)
a) pollen grains b) root tips
c) young leaves d) endosperm.

UNIT IX - Plant Ecology

1. Plants which produce characteristic pneumatophores and show vivipary belong to (NEET 2017)
a) **Halophytes** b) psammophytes
c) hydrophytes d) mesophytes

2. Mycorrhizae are the example of (NEET I 2017)
a) amensalism b) antibiosis
c) **mutualism** d) fungistatis

3. If '+' sign is assigned to beneficial interaction, '-' sign to detrimental and '0' sign to neutral interaction, then the population interaction represented by '+' '-' refers to (NEET 2016)
a) mutualism b) amensalism
c) commensalism d) **parasitism**

4. Which of the following is correctly matched? (NEET Phase 2 – 2016)
a) Aerenchyma - *Opuntia*
b) Age pyramid - Biome
c) ***Parthenium hysterophorus*** - **Threat to biodiversity**
d) Stratification - Population

5. An association of individuals of different species living in the same habitat and having functional interactions is (Re-AIPMT 2015)
a) Population b) Ecological niche
c) **Biotic community** d) Ecosystem

6. Roots play in significant role in absorption of water in (Re-AIPMT 2015)
a) Wheat b) Sunflower
c) ***Pistia*** d) Pea

7. If we uncover half of the forest covering the earth, what crisis will be produced at most and the first? (AIPMT 1996)
a. Some species will be extinct
b. Population and ecological imbalance will rise up
c. Energy crisis will occur
d. Rest half forests will maintain this imbalance.

8. Most animals are tree dwellers in a (AIPMT 2015)
a) **Tropical rain forest**
b)Coniferous forest
c) Thorn woodland
d) Temperate deciduous fo

9. *Cuscuta* is an example of (AIPMT Mains 2012)
a) **Ectoparasitism** b) Brood parasitism
c) Predation d) Endoparasitism

10. Large woody vines are more commonly found in (AIPMT Prelims 2011)
a) Alphine forests
b) Temperate forests
c) Mangroves
d) **Tropical rain forests**

11. Niche overlap indicates (AIPMT Prelims 2006)
a) Active co-operation between two species

TN_GOVT_BOTANY_XII_PAGES_216-279 CH 10.indd 272 28-12-2022 15:39:23