8. Show that $y = a\cos bx$ is a solution of the differential equation $\frac{d^2y}{dx^2} + b^2y = 0$.

Now, we discuss some standard methods of solving certain type of differential equations of the first order and first degree.

10.6 Solution of First Order and First Degree Differential Equations

10.6.1 Variables Separable Method

In solving differential equations, separation of variables was introduced initially by Leibniz and later it was formulated by John Bernoulli in the year 1694.

A first order differential equation is separable if it can be written as $h(y)y' = g(x)$ where the left side is a product of y' and a function of y and the right side is a function of x. Rewriting a separable differential equation in this form is called the method of separation of variables.

Finding a solution to a first order differential equation will be simple if the variables in the equation can be separated. An equation of the form $f_1(x)g_1(y)dx + f_2(x)g_2(y)dy = 0$ is called an equation with **variable separable** or simply a **separable equation**.

Rewrite the given differential equation as $\frac{f_1(x)}{f_2(x)}dx = -\frac{g_2(y)}{g_1(y)}dy$. ...(1)

Integration of both sides of (1) yields the general solution of the given differential equation as

$\int\frac{f_1(x)}{f_2(x)}dx = -\int\frac{g_2(y)}{g_1(y)}dy + C$, where C is an arbitrary constant.

Remarks

1. No need to add arbitrary constants on both sides as the two arbitrary constants are combined together as a single arbitrary constant.
2. A solution with this arbitrary constant is the general solution of the differential equation.

"Solving a differential equation" is also referred to as "integrating a differential equation", since the process of finding the solution to a differential equation involves integration.

Example 10.11

Solve $\left(1+x^2\right)\frac{dy}{dx} = 1+y^2$.



Solution

Given that $\left(1+x^2\right)\frac{dy}{dx} = 1+y^2$. ... (1)

The given equation is written in the variables separable form

$$\frac{dy}{1+y^2} = \frac{dx}{1+x^2}.$$ ... (2)

Integrating both sides of (2), we get $\tan^{-1}y = \tan^{-1}x + C$. ... (3)

But $\tan^{-1}y - \tan^{-1}x = \tan^{-1}\left(\frac{y-x}{1+xy}\right)$. ... (4)

12th_Maths_Vol 2_EM_CH 10_Differential Equation.indd 158 16-12-2022 16:47:39