10. On a multiple-choice exam with 3 possible destructives for each of the 5 questions, the probability that a student will get 4 or more correct answers just by guessing is

(1) $\frac{11}{243}$ (2) $\frac{3}{8}$ (3) $\frac{1}{243}$ (4) $\frac{5}{243}$

11. If $P(X=0) = 1 - P(X=1)$. If $E(X) = 3\text{Var}(X)$, then $P(X=0)$ is

(1) $\frac{2}{3}$ (2) $\frac{2}{5}$ (3) $\frac{1}{5}$ (4) $\frac{1}{3}$

12. If X is a binomial random variable with expected value 6 and variance 2.4, then $P(X=5)$ is

(1) $\binom{10}{5}\left(\frac{3}{5}\right)^6\left(\frac{2}{5}\right)^4$ (2) $\binom{10}{5}\left(\frac{3}{5}\right)^{10}$

(3) $\binom{10}{5}\left(\frac{3}{5}\right)^4\left(\frac{2}{5}\right)^6$ (4) $\binom{10}{5}\left(\frac{3}{5}\right)^5\left(\frac{2}{5}\right)^5$

13. The random variable X has the probability density function

$$f(x) = \begin{cases} ax+b & 0<x<1 \\ 0 & \text{otherwise} \end{cases}$$

and $E(X) = \frac{7}{12}$, then a and b are respectively

(1) 1 and $\frac{1}{2}$ (2) $\frac{1}{2}$ and 1 (3) 2 and 1 (4) 1 and 2

14. Suppose that X takes on one of the values 0, 1, and 2. If for some constant k, $P(X=i) = k\,P(X=i-1)$ for $i=1,2$ and $P(X=0) = \frac{1}{7}$, then the value of k is

(1) 1 (2) 2 (3) 3 (4) 4

15. Which of the following is a discrete random variable?
I. The number of cars crossing a particular signal in a day.
II. The number of customers in a queue to buy train tickets at a moment.
III. The time taken to complete a telephone call.

(1) I and II (2) II only (3) III only (4) II and III

16. If $f(x) = \begin{cases} 2x & 0 \le x \le a \\ 0 & \text{otherwise} \end{cases}$ is a probability density function of a random variable, then the value of a is

(1) 1 (2) 2 (3) 3 (4) 4

17. The probability mass function of a random variable is defined as:

x	–2	–1	0	1	2
$f(x)$	k	$2k$	$3k$	$4k$	$5k$

Then $E(X)$ is equal to:

(1) $\frac{1}{15}$ (2) $\frac{1}{10}$ (3) $\frac{1}{3}$ (4) $\frac{2}{3}$

12th_Maths_Vol 2_EM_CH 11_Probability Distributions_29-05-20.indd 220 16-12-2022 16:52:51