


Figure 3.48 Toroid – Amperian loop

(a) Open space interior to the toroid

Let us calculate the magnetic field B_P at point P. We construct an Amperian loop 1 of radius r_1 around the point P as shown in Figure 3.48. For simplicity, we take circular loop so that the length of the loop is its circumference.

$$L_1 = 2\pi r_1$$

Ampère's circuital law for the loop 1 is

$$\oint_{loop1} \vec{B}_P . \vec{dl} = \mu_\circ I_{enclosed}$$

Since, the loop1 encloses no current, $I_{enclosed} = 0$

$$\oint_{loop1} \vec{B}_P . \vec{dl} = 0$$

This is possible only if the magnetic field at point P vanishes i.e.

$$\vec{B}_P = 0$$

(b) Open space exterior to the toroid

Let us calculate the magnetic field B_Q at point Q. We construct an Amperian loop 3 of radius r_3 around the point Q as shown in Figure 3.48. The length of the loop is

$$L_3 = 2\pi r_3$$

Ampère's circuital law for the loop 3 is

$$\oint_{loop3} \vec{B}_Q . \vec{dl} = \mu_\circ I_{enclosed}$$

Since, in each turn of the toroid loop, current coming out of the plane of paper is cancelled by the current going into the plane of paper. Thus, $I_{enclosed} = 0$

$$\oint_{loop3} \vec{B}_Q . \vec{dl} = 0$$

This is possible only if the magnetic field at point Q vanishes i.e.

$$\vec{B}_Q = 0$$

(c) Inside the toroid

Let us calculate the magnetic field B_S at point S by constructing an Amperian loop 2 of radius r_2 around the point S as shown in Figure 3.48. The length of the loop is

$$L_2 = 2\pi r_2$$

Ampere's circuital law for the loop 2 is

$$\oint_{loop2} \vec{B}_S . \vec{dl} = \mu_\circ I_{enclosed}$$

Let I be the current passing through the toroid and N be the number of turns of the toroid, then

$$I_{enclosed} = NI$$

and $$\oint_{loop2} \vec{B}_S . \vec{dl} = \oint_{loop2} B dl \cos\theta = B 2\pi r_2$$

$$\oint_{loop2} \vec{B}_S . \vec{dl} = \mu_\circ NI$$

UNIT-3(XII-Physics_Vol-1).indd 174 04-03-2019 10:34:53