EXERCISE 6.9

1. Find the equation of the plane passing through the line of intersection of the planes $\vec{r}\cdot\left(2\hat{i}-7\hat{j}+4\hat{k}\right)=3$ and $3x-5y+4z+11=0$, and the point $(-2,1,3)$.
2. Find the equation of the plane passing through the line of intersection of the planes $x+2y+3z=2$ and $x-y+z=3$, and at a distance $\frac{2}{\sqrt{3}}$ from the point $(3,1,-1)$.
3. Find the angle between the line $\vec{r}=\left(2\hat{i}-\hat{j}+\hat{k}\right)+t\left(\hat{i}+2\hat{j}-2\hat{k}\right)$ and the plane $\vec{r}\cdot\left(6\hat{i}+3\hat{j}+2\hat{k}\right)=8$
4. Find the angle between the planes $\vec{r}\cdot\left(\hat{i}+\hat{j}-2\hat{k}\right)=3$ and $2x-2y+z=2$.
5. Find the equation of the plane which passes through the point $(3,4,-1)$ and is parallel to the plane $2x-3y+5z+7=0$. Also, find the distance between the two planes.
6. Find the length of the perpendicular from the point $(1,-2,3)$ to the plane $x-y+z=5$.
7. Find the point of intersection of the line $x-1=\frac{y}{2}=z+1$ with the plane $2x-y+2z=2$. Also, find the angle between the line and the plane.
8. Find the coordinates of the foot of the perpendicular and length of the perpendicular from the point (4,3,2) to the plane $x+2y+3z=2$.

EXERCISE 6.10

Choose the correct or the most suitable answer from the given four alternatives :

1. If $\vec{a}$ and $\vec{b}$ are parallel vectors, then $[\vec{a},\vec{c},\vec{b}]$ is equal to
   (1) 2 (2) –1 (3) 1 (4) 0
2. If a vector $\vec{\alpha}$ lies in the plane of $\vec{\beta}$ and $\vec{\gamma}$, then
   (1) $[\vec{\alpha},\vec{\beta},\vec{\gamma}]=1$ (2) $[\vec{\alpha},\vec{\beta},\vec{\gamma}]=-1$ (3) $[\vec{\alpha},\vec{\beta},\vec{\gamma}]=0$ (4) $[\vec{\alpha},\vec{\beta},\vec{\gamma}]=2$
3. If $\vec{a}\cdot\vec{b}=\vec{b}\cdot\vec{c}=\vec{c}\cdot\vec{a}=0$, then the value of $[\vec{a},\vec{b},\vec{c}]$ is
   (1) $|\vec{a}|\,|\vec{b}|\,|\vec{c}|$ (2) $\frac{1}{3}|\vec{a}|\,|\vec{b}|\,|\vec{c}|$ (3) 1 (4) –1
4. If $\vec{a},\vec{b},\vec{c}$ are three unit vectors such that $\vec{a}$ is perpendicular to $\vec{b}$, and is parallel to $\vec{c}$ then $\vec{a}\times(\vec{b}\times\vec{c})$ is equal to
   (1) $\vec{a}$ (2) $\vec{b}$ (3) $\vec{c}$ (4) $\vec{0}$
5. If $[\vec{a},\vec{b},\vec{c}]=1$, then the value of $\frac{\vec{a}\cdot(\vec{b}\times\vec{c})}{(\vec{c}\times\vec{a})\cdot\vec{b}}+\frac{\vec{b}\cdot(\vec{c}\times\vec{a})}{(\vec{a}\times\vec{b})\cdot\vec{c}}+\frac{\vec{c}\cdot(\vec{a}\times\vec{b})}{(\vec{c}\times\vec{b})\cdot\vec{a}}$ is
   (1) 1 (2) –1 (3) 2 (4) 3
6. The volume of the parallelepiped with its edges represented by the vectors $\hat{i}+\hat{j}$, $\hat{i}+2\hat{j}$, $\hat{i}+\hat{j}+\pi\hat{k}$ is
   (1) $\frac{\pi}{2}$ (2) $\frac{\pi}{3}$ (3) π (4) $\frac{\pi}{4}$

12th_Maths_EM_Vol1_CH 6 Vector Algebra.indd 276 16-12-2022 14:25:59