selection pressure to the black coloured moths. The above proof shows that in a population, organisms that can adapt will survive and produce more progenies resulting in increase in population through natural selection.

Artificial selection is a byproduct of human exploitation of forests, oceans and fisheries or the use of pesticides, herbicides or drugs. For hundreds of years humans have selected various types of dogs, all of which are variants of the single species of dog. If human beings can produce new varieties in short period, then **"nature"** with its vast resources and long duration can easily produce new species by selection.

6.5.6 Adaptive Radiation

The evolutionary process which produces new species diverged from a single ancestral form becomes adapted to newly invaded habitats is called adaptive radiation. Adaptive radiations are best exemplified in closely related groups that have evolved in relatively short time. Darwin's finches and Australian marsupials are best examples for adaptive radiation. When more than one adaptive radiation occurs in an isolated geographical area, having the same structural and functional similarity it is due to convergent evolution.

Darwin's finches

Their common ancestor arrived on the Galapagos about 2 million years ago. During that time, Darwin's finches have evolved into 14 recognized species differing in body size, beak shape and feeding behavior. Changes in the size and form of the beak have enabled different species to utilize different food resources such as insects, seeds, nectar from cactus flowers and blood from iguanas, all driven by Natural selection. **Fig. 6.5** represents some of the finches observed by Darwin. Genetic variation in the ALX1 gene in the DNA of Darwin finches is associated with variation in the beak shape. Mild mutation in the ALX1 gene leads to phenotypic change in the shape of the beak of the Darwin finches.

Marsupials in Australia and placental mammals in North America are two subclasses of mammals they have adapted in similar way to a particular food resource, locomotory skill or climate. They were separated from the common ancestor more than 100 million year ago and each lineage continued to evolve independently. Despite temporal and geographical separation, marsupials in Australia and placental mammals in North America have produced varieties of species living in similar habitats with similar ways of life. Their overall resemblance in shape, locomotory mode, feeding and foraging are superimposed upon different modes of reproduction. This feature reflects their distinctive evolutionary relationships.

Over 200 species of marsupials live in Australia along with many fewer species of placental mammals. The marsupials have undergone adaptive radiation to occupy the diverse habitats in Australia, just as the placental mammals have radiated across North America.




Fig 6.5 Darwin's finches

XII Std Zoology Chapter 6 EM.indd 102 23/12/2022 16:12:50