This increases the emission of CO_2, a GHG, leading to climate change. Due to large scale deforestation, the emitted CO_2 cannot be absorbed fully, and its concentration in the air increases. Climate change increases land and ocean temperature, changes precipitation patterns and raises the sea level. This inturn results in melting of glaciers, water inundation, less predictability of weather patterns, extreme weather conditions, outbreak of squalor diseases, migration of animals and loss of trees in forest. Thus, climate change is an imminent danger to the existing biodiversity **(Fig. 12.4)**.




Fig. 12.4 Links between biodiversity, climate change and human well - being

Shifting or Jhum cultivation (Slash-and-burn agriculture)

In shifting cultivation, plots of natural tree vegetation are burnt away and the cleared patches are farmed for 2-3 seasons, after which their fertility reduces to a point where crop production is no longer profitable. The farmer then abandons this patch and cuts down a new patch of forest trees elsewhere for crop production. This system is practiced in north-eastern regions of India. When vast areas are cleared and burnt, it results in loss of forest cover, pollution and discharge of CO_2 which in turn attributes to loss of habitat and climate change which has an impact on the faunal diversity of that regions.

Coextinctions

Coextinction of a species is the loss of a species as a consequence of the extinction of another. (Eg., orchid bees and forest trees by cross pollination). Extinction of one will automatically cause extinction of the other. Another example for co-extinction is the connection between Calvaria tree and the extinct bird of Mauritius Island, the Dodo. The Calvaria tree is dependent on the Dodo bird for completion of its life cycle. The mutualistic association is that the tough horny endocarp of the seeds of Calvaria tree are made permeable by the actions of the large stones in bird's gizzard and digestive juices thereby facilitating easier germination. The extinction of the Dodo bird led to the imminent danger of the Calvaria tree coextinction.

Intensive agriculture

Spread of agriculture is sometimes at the cost of wetlands, grasslands and forests. Intensive agriculture is based on a few high yielding varieties. As a result, there is reduction in the genetic diversity. It also increases vulnerability of the crop plants to sudden attack by pathogens and pests. There are only few varieties of traditional paddy strains today due to use to hybrid varieties in Tamil Nadu.

12.5.1 Loss of Biodiversity

Species have been evolving and dying out (extinction) ever since the origin of life. However, species are now becoming extinct at a faster rate. This is destabilizing the ecological stability and the distribution of biological diversity on earth. Human activities greatly contribute to the loss of biodiversity. Natural resources such as land, water and organisms are indiscriminately exploited by human beings.

According to the Convention of Biological Diversity, direct and indirect human activities have a detrimental effect

XII Std Zoology Chapter 12 EM.indd 207 23/12/2022 16:52:09