

Simulation of the unstable rotation of a cuboid

Books in space

Ivar Postma Eamon Nerbonne

Introduction to Computational Science
School for Computing and Cognition
University of Groningen

December 1, 2011

Experiment in space



- ▶ <http://www.youtube.com/watch?v=GgVp0orcKqc>

Stable and unstable rotation

- ▶ The book can stably rotate around two of its axis
- ▶ Rotation around the third axis is not stable

Modelling the behaviour

We make a few simplifications

- ▶ The book is a cuboid
- ▶ The density is constant

Model

State vector

For a point on the body we keep track of the position, the orientation, the total linear momentum and the total angular momentum

$$S(t) = \begin{pmatrix} x(t) \\ R(t) \\ P(t) \\ L(t) \end{pmatrix}$$

Model

Updating the state vector

To update we need the velocity, the angular speed, the force and the torque (moment of force)

$$\frac{d}{dt}S(t) = \begin{pmatrix} v(t) \\ \omega(t) * R(t) \\ F(t) \\ \tau(t) \end{pmatrix}$$

Model

More simplifications

When a book is rotating in space, the model is simple because

- ▶ there is no force, $F(t) = 0$
- ▶ there is no torque, $\tau(t) = 0$

Thus, we only need to calculate the velocity and angular speed

Model

Updating the state vector

Velocity is given by

$$v(t) = \omega(t)r(t)$$

Angular speed is given by

$$\omega(t) = I(t)^{-1}L(t)$$

$I(t)$ depends on $R(t)$ and the moment of inertia of the body I_{body}

$$I(t) = R(t)I_{body}R(t)^T$$

Moment of inertia

- ▶ The moment of inertia of an object is a measure of the resistance of an object to changes in the rotation
- ▶ For a cuboid the moment of inertia is given by

$$I_{body} = \frac{1}{12} \begin{pmatrix} M(b^2 + c^2) & 0 & 0 \\ 0 & M(a^2 + c^2) & 0 \\ 0 & 0 & M(a^2 + b^2) \end{pmatrix}$$

where a , b and c are the dimensions of the cuboid

Methods to solve the calculations

- ▶ Euler intergration
- ▶ Midpoint method
- ▶ 4th order Runga Kutta

Implementation

We intend to implement this in C/C++ using OpenGL for the visualisation

Questions?