

Simulation of the unstable rotation of a cuboid

Books in space

Ivar Postma Eamon Nerbonne

Introduction to Computational Science
School for Computing and Cognition
University of Groningen

January 19, 2011

What you should remember from last time...

- ▶ Rotation of a rigid body; a book
- ▶ Clip of an experiment in space
- ▶ Rotation around two of the main axes is stable, rotation around the third is not

String particle simulation

- ▶ Lecture on Molecular Dynamics: simulation of vibrating string with particles
- ▶ Springs create interacting forces between particles
- ▶ From forces, calculate update velocity of points ($a = \frac{F}{m}$)
- ▶ From velocity, calculate update position of points

Particle simulation


Model of a book

- ▶ Eight particles, located at the corners
- ▶ Each point has the same mass
- ▶ Interacting force (spring) between each pair of particles
- ▶ Force is proportional to distance between the particles minus initial distance

Particle simulation

Model of a book

- ▶ Eight particles, located at the corners
- ▶ Each point has the same mass
- ▶ Interacting force (spring) between each pair of particles
- ▶ Force is proportional to distance between the particles minus initial distance



Particle simulation

Computations

- ▶ Calculate force on particle i :

$$F_i(t) = \sum_j (x_i(t) - x_j(t)) - (x_i(0) - x_j(0))$$

- ▶ Calculate velocity of particle i :

$$v_i(t + \Delta t) = v_i(t) + F_i(t) \times \Delta t$$

- ▶ Calculate position of particle i :

$$x_i(t + \Delta t) = x_i(t) + v_i(t) \times \Delta t$$

Particle simulation


Computations

Using Euler integration, for n particles, in each time step we need to compute

- ▶ $\frac{(n-1)^2}{2}$ forces
- ▶ n velocities
- ▶ n positions

Particle simulation

Demonstration



Particle simulation

Observations

- ▶ Book behaves naturally
- ▶ Rotation is stable around two axis and unstable around the third
- ▶ After some time the simulation “explodes”


Particle simulation

Normalisation

- ▶ Euler integration introduces a small error in each time step
- ▶ Shape of book changes eventually
- ▶ Other techniques can reduce, but not eliminate the error
- ▶ Normalise the model using the conservation of energy

Particle simulation

Demonstration with normalisation



Particle simulation

Normalisation

- ▶ Energy is conserved
- ▶ Angular momentum is transformed into vibration

A different approach

- ▶ Different approach introduces last time
- ▶ Calculate angular velocity to update the orientation

Angular velocity based simulation

Computations

- ▶ Angular velocity calculated from angular momentum and moment of inertia

$$I(t) = R(t) \times I_{body} \times R(t)^T$$

$$\omega(t) = I(t)^{-1} \times L$$

- ▶ Change in orientation calculated from angular velocity

$$R(t + \Delta t) = R(t) + R(t) \times \omega^*(t) \times \Delta t$$

Angular velocity based simulation


Computations

Using Euler integration, in each time step we need to do

- ▶ 5 matrix multiplications
- ▶ 1 vector to tensor calculation
- ▶ 1 matrix inverse

Angular velocity based simulation

Demonstration




Calculating the orientation

Normalisation

- ▶ Euler integration introduces a small error in each time step
- ▶ Book changes shape
- ▶ Orientation matrix normalised by singular value decomposition (SVD)

Angular velocity based simulation

Demonstration with normalisation




Quaternions

- ▶ Orientation has 3 degrees of freedom, but a 3×3 matrix (9 degrees of freedom)
- ▶ Orientation with quaternions has 4 degrees of freedom
- ▶ Again, normalisation to prevent shape from changing

Angular velocity based simulation

Demonstration with quaternions



Conclusion

- ▶ (Un)stable rotation of a book can be simulated with particles
- ▶ Simulation eventually produces “unnatural” results
- ▶ Other approach preserves angular momentum
- ▶ Simulation is more robust
- ▶ Only works when moment of inertia is known

Questions?

?

Questions

about particle simulation

Q: What happens if you change the spring constant in calculating the forces?

A: For a high spring constant the shape and size of the book is contained better, but the book quickly goes to the vibrating state. For a low spring constant it takes longer to reach the vibrating state, but the shape and size is distorted more.

Q: Why don't you normalise the distance between particles?

A: If we do that, how will we calculate the forces?

Questions

about particle simulation

Q: Can you improve this model?

A: There are several extensions one can think of. For example, damping may produce better results.

Questions

about quaternions

Q: What are quaternions?

A: They're an extension to complex numbers of the form $q = a + b \cdot i + c \cdot j + d \cdot k$ that can be used to describe a rotation in 3d space. For a point at position $p(0, x, y, z)$ the position after rotation is $p'(0, x', y', z') = q \cdot p \cdot \bar{q}$.

Questions

about quaternions vs. matrix method

Q: Which method is faster?

A: In theory, quaternions produce better results and can be normalised less frequently. However, in this case the simulation is fast enough to do a normalisation in every time step.