



CS215 DISCRETE MATH

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: wangqi@sustech.edu.cn

Recursion

- Recursive computer programs or algorithms often lead to inductive analysis.



Recursion

- Recursive computer programs or algorithms often lead to *inductive analysis*.
- A classical example of *recursion* is the **Towers of Hanoi** Problem.

Towers of Hanoi



3 - 1



Towers of Hanoi



- 3 pegs; n disks of different sizes
- A *legal move* takes a disk from one peg and moves it onto another peg so that **it is not on top of a smaller disk**
- **Problem:** Find an (efficient) way to move all of the disks from one peg to another


Towers of Hanoi



Towers of Hanoi




legal move




Towers of Hanoi



legal move




legal move




Towers of Hanoi




legal move



legal move




not legal




Towers of Hanoi




legal move



legal move



not legal



legal move



Towers of Hanoi

- **Problem:** Start with n disks on leftmost peg




Towers of Hanoi

- **Problem:** Start with n disks on leftmost peg using only legal moves



Towers of Hanoi

- **Problem:** Start with n disks on leftmost peg using only legal moves
move all disks to rightmost peg.



Towers of Hanoi

- **Problem:** Start with n disks on leftmost peg
using only legal moves
move all disks to rightmost peg.



Given $i, j \in \{1, 2, 3\}$, let
 $\overline{\{i, j\}} = \{1, 2, 3\} - \{i\} - \{j\}$,
i.e., $\overline{\{1, 2\}} = \{3\}$, $\overline{\{1, 3\}} = \{2\}$,
 $\overline{\{2, 3\}} = \{1\}$.



Towers of Hanoi

- General solution



Towers of Hanoi

- General solution

Recursion Base:

If $n = 1$, moving one disk from i to j is easy. Just move it.



Towers of Hanoi

- General solution

Recursion Base:

If $n = 1$, moving one disk from i to j is easy. Just move it.



Towers of Hanoi



To move $n > 1$ disks from i to j

Towers of Hanoi



To move $n > 1$ disks from i to j

1)



move top $n - 1$ disks from i to $\overline{\{i, j\}}$


Towers of Hanoi



1)



2)



To move $n > 1$ disks from i to j

move top $n - 1$ disks from i to $\overline{\{i, j\}}$

move **largest** disk from i to j

Towers of Hanoi



1)



2)



3)



To move $n > 1$ disks from i to j

move top $n - 1$ disks from i to $\overline{\{i, j\}}$

move largest disk from i to j

move top $n - 1$ disks from $\overline{\{i, j\}}$ to j

Towers of Hanoi

```
3 public class Hanoi
4 {
5
6     public void move(int n, char a, char b, char c)
7     {
8         if (n == 1)
9             System.out.println("plate " + n + " from " + a + " to " + c);
10        else
11        {
12            move(n-1,a,c,b);
13            System.out.println("plate " + n + " from " + a + " to " + c);
14            move(n-1,b,a,c);
15        }
16    }
17 }
18
```

Towers of Hanoi

To move n disks from i to j

i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$

ii) move largest disk from i to j

iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j

Towers of Hanoi

- To prove **correctness** of solution, we are implicitly using **induction**

To move n disks from i to j

- move top $n - 1$ disks from i to $\overline{\{i, j\}}$
- move largest disk from i to j
- move top $n - 1$ disks from $\overline{\{i, j\}}$ to j



Towers of Hanoi

- To prove **correctness** of solution, we are implicitly using **induction**
- $p(n)$ is statement that algorithm is correct for n

To move n disks from i to j

- i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$
- ii) move largest disk from i to j
- iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j



Towers of Hanoi

- To prove **correctness** of solution, we are implicitly using **induction**

- $p(n)$ is statement that algorithm is correct for n

- $p(1)$ is statement that algorithm works for $n = 1$ disks, which is obviously true

To move n disks from i to j

i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$

ii) move largest disk from i to j

iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j

Towers of Hanoi

- To prove **correctness** of solution, we are implicitly using **induction**

- $p(n)$ is statement that algorithm is correct for n

- $p(1)$ is statement that algorithm works for $n = 1$ disks, which is obviously true

- $p(n - 1) \rightarrow p(n)$ is **recursion** statement that

if our algorithm works for $n - 1$ disks, then we can build a correct solution for n disks

To move n disks from i to j

i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$

ii) move largest disk from i to j

iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j

Towers of Hanoi

■ Running time

$M(n)$ is number of disk moves needed for n disks

To move n disks from i to j

- i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$
- ii) move largest disk from i to j
- iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j

Towers of Hanoi

■ Running time

$M(n)$ is number of disk moves needed for n disks

To move n disks from i to j

- i) move top $n - 1$ disks from i to $\overline{\{i, j\}}$
- ii) move largest disk from i to j
- iii) move top $n - 1$ disks from $\overline{\{i, j\}}$ to j

$$M(1) = 1$$

$$\text{if } n > 1, \text{ then } M(n) = 2M(n - 1) + 1$$

Towers of Hanoi

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$



Towers of Hanoi

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$

- Iterating the recurrence gives

$$M(1) = 1, M(2) = 3, M(3) = 7, \\ M(4) = 15, M(5) = 31, \dots$$



Towers of Hanoi

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$

- Iterating the recurrence gives

$$M(1) = 1, M(2) = 3, M(3) = 7, \\ M(4) = 15, M(5) = 31, \dots$$

- We *guess* that $M(n) = 2^n - 1$



Towers of Hanoi

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$

- Iterating the recurrence gives

$$M(1) = 1, M(2) = 3, M(3) = 7, \\ M(4) = 15, M(5) = 31, \dots$$

- We *guess* that $M(n) = 2^n - 1$
We'll prove this by induction



Towers of Hanoi

- We saw that $M(1) = 1$ and that
- $M(n) = 2M(n - 1) + 1$ for $n > 1$

- Iterating the recurrence gives

$$M(1) = 1, M(2) = 3, M(3) = 7, \\ M(4) = 15, M(5) = 31, \dots$$

- We *guess* that $M(n) = 2^n - 1$

We'll prove this *by induction*

Later, we'll also see how to solve *without guessing*



Towers of Hanoi

- Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$

We show that $M(n) = 2^n - 1$.

Towers of Hanoi

- Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$

We show that $M(n) = 2^n - 1$.

Proof. (by induction)

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

For the inductive step, assume that $M(n-1) = 2^{n-1} - 1$ for $n > 1$.



Towers of Hanoi

- Formally, given

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$

We show that $M(n) = 2^n - 1$.

Proof. (by induction)

The base case $n = 1$ is true, since $2^1 - 1 = 1$.

For the inductive step, assume that $M(n-1) = 2^{n-1} - 1$ for $n > 1$.

Then $M(n) = 2M(n-1) + 1 = 2(2^{n-1} - 1) + 1 = 2^n - 1$



Towers of Hanoi

- Note that we used induction twice.



Towers of Hanoi

- Note that we used **induction twice**.
- The first time was to **derive** correctness of algorithm and the recurrence

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$



Towers of Hanoi

- Note that we used **induction twice**.
- The first time was to **derive** correctness of algorithm and the recurrence

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n-1) + 1 & \text{otherwise} \end{cases}$$

- The second time was to **derive** the **closed form solution** $M(n) = 2^n - 1$ of the recurrence.



Recurrences

- A *recurrence equation* or *recurrence* for a function defined on the set of integers $\geq b$ is one that tells us how to compute the n th value from some or all the first $n - 1$ values.



Recurrences

- A *recurrence equation* or *recurrence* for a function defined on the set of integers $\geq b$ is one that tells us **how to compute the n th value from some or all the first $n - 1$ values.**

To completely specify a function on the basis of a recurrence, we have to give the *initial condition(s)* (a.k.a. the *base case(s)*) for the recurrence.



Recurrences

- A *recurrence equation* or *recurrence* for a function defined on the set of integers $\geq b$ is one that tells us **how to compute the n th value from some or all the first $n - 1$ values.**

To completely specify a function on the basis of a recurrence, we have to give the *initial condition(s)* (a.k.a. the *base case(s)*) for the recurrence.

$$M(n) = \begin{cases} 1 & \text{if } n = 1 \\ 2M(n - 1) + 1 & \text{otherwise} \end{cases} \quad \text{Towers of Hanoi}$$

$$F(n) = \begin{cases} 1 & \text{if } n = 0, 1 \\ F(n - 1) + F(n - 2) & \text{otherwise} \end{cases} \quad \text{Fibonacci Sequence}$$



Recurrences

- **Example 2:** Let $S(n)$ be the number of subsets of a set of size n . What is the formula for $S(n)$?

The empty set, of size $n = 0$ has only one subset (itself), so $S(0) = 1$.

It is not difficult to see that

$$S(1) = 2, S(2) = 4, S(3) = 8$$



Recurrences

- **Example 2:** Let $S(n)$ be the number of subsets of a set of size n . What is the formula for $S(n)$?

The empty set, of size $n = 0$ has only one subset (itself), so $S(0) = 1$.

It is not difficult to see that

$$S(1) = 2, S(2) = 4, S(3) = 8$$

We “guess” that $S(n) = 2^n$. But, in order to prove formula, we’ll need to think recursively.



Recurrences

- Consider the eight subsets of $\{1, 2, 3\}$:
 $\emptyset, \{1\}, \{2\}, \{1, 2\}, \{3\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}$



Recurrences

- Consider the eight subsets of $\{1, 2, 3\}$:

$\emptyset, \{1\}, \{2\}, \{1, 2\}, \{3\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}$

\emptyset	$\{1\}$	$\{2\}$	$\{1, 2\}$
$\{3\}$	$\{1, 3\}$	$\{2, 3\}$	$\{1, 2, 3\}$



Recurrences

- Consider the eight subsets of $\{1, 2, 3\}$:

$\emptyset, \{1\}, \{2\}, \{1, 2\}, \{3\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}$

\emptyset	$\{1\}$	$\{2\}$	$\{1, 2\}$
$\{3\}$	$\{1, 3\}$	$\{2, 3\}$	$\{1, 2, 3\}$

First four subsets are exactly the subsets of $\{1, 2\}$, while second four are the subsets of $\{1, 2\}$ with 3 added into each.

Recurrences

- Consider the eight subsets of $\{1, 2, 3\}$:

$\emptyset, \{1\}, \{2\}, \{1, 2\}, \{3\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}$

\emptyset	$\{1\}$	$\{2\}$	$\{1, 2\}$
$\{3\}$	$\{1, 3\}$	$\{2, 3\}$	$\{1, 2, 3\}$

First four subsets are exactly the subsets of $\{1, 2\}$, while second four are the subsets of $\{1, 2\}$ with 3 added into each.

So, we get a subset of $\{1, 2, 3\}$ either by taking a subset of $\{1, 2\}$ or by adjoining 3 to a subset of $\{1, 2\}$.



Recurrences

- Consider the eight subsets of $\{1, 2, 3\}$:

$\emptyset, \{1\}, \{2\}, \{1, 2\}, \{3\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}$

\emptyset	$\{1\}$	$\{2\}$	$\{1, 2\}$
$\{3\}$	$\{1, 3\}$	$\{2, 3\}$	$\{1, 2, 3\}$

First four subsets are exactly the subsets of $\{1, 2\}$, while second four are the subsets of $\{1, 2\}$ with 3 added into each.

So, we get a subset of $\{1, 2, 3\}$ either by taking a subset of $\{1, 2\}$ or by adjoining 3 to a subset of $\{1, 2\}$.

This suggests that the recurrence for the number of subsets of an n -element set $\{1, 2, \dots, n\}$ is

$$S(n) = \begin{cases} 1 & \text{if } n = 0 \\ 2S(n-1) & \text{if } n \geq 1 \end{cases}$$



Recurrences

- **Proof.** of correctness of this recurrence



Recurrences

- **Proof.** of correctness of this recurrence

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain the element n .



Recurrences

■ **Proof.** of correctness of this recurrence

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain the element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .



Recurrences

■ **Proof.** of correctness of this recurrence

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain the element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .

So, the number of subsets containing n is exactly the same as the number of subsets not containing n .



Recurrences

■ **Proof.** of correctness of this recurrence

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain the element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .

So, the number of subsets containing n is exactly the same as the number of subsets not containing n .

Thus, if $n > 1$, then $S(n) = 2S(n - 1)$.



Recurrences

■ **Proof.** of correctness of this recurrence

The subsets of $\{1, 2, \dots, n\}$ can be partitioned according to whether or not they contain the element n .

Each subset S containing n can be constructed in a unique fashion by adding n to the subset $S - \{n\}$ not containing n .

Each subset S not containing n can be constructed by removing n from the unique set $S \cup \{n\}$ containing n .

So, the number of subsets containing n is exactly the same as the number of subsets not containing n .

Thus, if $n > 1$, then $S(n) = 2S(n - 1)$.

Proof by induction is easy.



Iterating a Recurrence

- Let $T(n) = rT(n-1) + a$,
where r and a are constants.



Iterating a Recurrence

- Let $T(n) = rT(n-1) + a$,
where r and a are constants.

Find a **recurrence** that expresses

$T(n)$ in terms of $T(n-2)$

$T(n)$ in terms of $T(n-3)$

$T(n)$ in terms of $T(n-4)$

⋮

Iterating a Recurrence

- Let $T(n) = rT(n-1) + a$,
where r and a are constants.

Find a **recurrence** that expresses

$T(n)$ in terms of $T(n-2)$

$T(n)$ in terms of $T(n-3)$

$T(n)$ in terms of $T(n-4)$

⋮

Can we generalize this to find a **closed-form solution**?



Iterating a Recurrence

- Note that $T(n) = rT(n-1) + a$ implies that
 $\forall i < n, T(n-i) = rT((n-i)-1) + a$



Iterating a Recurrence

- Note that $T(n) = rT(n-1) + a$ implies that

$$\forall i < n, T(n-i) = rT((n-i)-1) + a$$

Then, we have

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2 T(n-2) + ra + a \\ &= r^2(rT(n-3) + a) + ra + a \\ &= r^3 T(n-3) + r^2 a + ra + a \\ &= r^3(rT(n-4) + a) + r^2 a + ra + a \\ &= r^4 T(n-4) + r^3 a + r^2 a + ra + a. \end{aligned}$$

Iterating a Recurrence

- Note that $T(n) = rT(n-1) + a$ implies that

$$\forall i < n, T(n-i) = rT((n-i)-1) + a$$

Then, we have

$$\begin{aligned} T(n) &= rT(n-1) + a \\ &= r(rT(n-2) + a) + a \\ &= r^2 T(n-2) + ra + a \\ &= r^2(rT(n-3) + a) + ra + a \\ &= r^3 T(n-3) + r^2 a + ra + a \\ &= r^3(rT(n-4) + a) + r^2 a + ra + a \\ &= r^4 T(n-4) + r^3 a + r^2 a + ra + a. \end{aligned}$$

Guess $T(n) = r^n T(0) + a \sum_{i=0}^{n-1} r^i$



Iterating a Recurrence

- The method we used to guess the solution is called *iterating the recurrence*, because we repeatedly (iteratively) use the recurrence.



Iterating a Recurrence

- The method we used to guess the solution is called *iterating the recurrence*, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “bottom-up” instead of “top-down”.



Iterating a Recurrence

- The method we used to guess the solution is called *iterating the recurrence*, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “*bottom-up*” instead of “*top-down*”.

$$T(0) = b$$

$$T(1) = rT(0) + a = rb + a$$

$$T(2) = rT(1) + a = r(rb + a) + a = r^2b + ra + a$$

$$T(3) = rT(2) + a = r^3b + r^2a + ra + a$$



Iterating a Recurrence

- The method we used to guess the solution is called *iterating the recurrence*, because we repeatedly (iteratively) use the recurrence.

Another approach is to iterate from the “bottom-up” instead of “top-down”.

$$T(0) = b$$

$$T(1) = rT(0) + a = rb + a$$

$$T(2) = rT(1) + a = r(rb + a) + a = r^2b + ra + a$$

$$T(3) = rT(2) + a = r^3b + r^2a + ra + a$$

This would lead to the same guess

$$T(n) = r^n b + a \sum_{i=0}^{n-1} r^i.$$



Formula of Recurrences

- **Theorem** If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .



Formula of Recurrences

- **Theorem** If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Proof by induction

The base case:

$$T(0) = r^0 b + a \frac{1 - r^0}{1 - r} = b.$$

So the formula is true when $n = 0$.

Now assume that $n > 0$ and

$$T(n-1) = r^{n-1} b + a \frac{1 - r^{n-1}}{1 - r}.$$



Formula of Recurrences

■ Proof by induction

$$\begin{aligned}T(n) &= rT(n-1) + a \\&= r \left(r^{n-1}b + a \frac{1-r^{n-1}}{1-r} \right) + a \\&= r^n b + \frac{ar - ar^n}{1-r} + a \\&= r^n b + \frac{ar - ar^n + a - ar}{1-r} \\&= r^n b + a \frac{1-r^n}{1-r}.\end{aligned}$$

Formula of Recurrences

- **Theorem** If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .



Formula of Recurrences

- **Theorem** If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Example:

$$T(n) = 3T(n-1) + 2 \text{ with } T(0) = 5$$



Formula of Recurrences

- **Theorem** If $T(n) = rT(n-1) + a$, $T(0) = b$, and $r \neq 1$, then

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

for all nonnegative integers n .

Example:

$$T(n) = 3T(n-1) + 2 \text{ with } T(0) = 5$$

Plugging $r = 3$, $a = 2$, $b = 5$ in the formula, gives

$$T(n) = 3^n \cdot 5 + 2 \frac{1 - 3^n}{1 - 3} = 3^n \cdot 6 - 1$$



First-Order Linear Recurrences

- A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a *first-order linear recurrence*.



First-Order Linear Recurrences

- A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a *first-order linear recurrence*.
 - ◇ **First order** because it only depends upon going back one step, i.e., $T(n-1)$



First-Order Linear Recurrences

- A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a *first-order linear recurrence*.

◇ **First order** because it only depends upon going back one step, i.e., $T(n-1)$

If it depends upon $T(n-2)$, it would be a **second-order** recurrence, e.g., $T(n) = T(n-1) + 2T(n-2)$.



First-Order Linear Recurrences

- A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a *first-order linear recurrence*.
 - ◇ **First order** because it only depends upon going back one step, i.e., $T(n-1)$

If it depends upon $T(n-2)$, it would be a **second-order** recurrence, e.g., $T(n) = T(n-1) + 2T(n-2)$.
 - ◇ **Linear** because $T(n-1)$ only appears to the **first power**.



First-Order Linear Recurrences

- A recurrence of the form $T(n) = f(n)T(n-1) + g(n)$ is called a *first-order linear recurrence*.
 - ◇ **First order** because it only depends upon going back one step, i.e., $T(n-1)$

If it depends upon $T(n-2)$, it would be a **second-order** recurrence, e.g., $T(n) = T(n-1) + 2T(n-2)$.
 - ◇ **Linear** because $T(n-1)$ only appears to the **first power**.

Something like $T(n) = (T(n-1))^2 + 3$ would be a **non-linear** first-order recurrence relation.



First-Order Linear Recurrences

- $T(n) = f(n)T(n-1) + g(n)$



First-Order Linear Recurrences

- $T(n) = f(n)T(n-1) + g(n)$

When $f(n)$ is a constant, say r , the general solution is almost as easy as we derived before. Iterating the recurrence gives

$$\begin{aligned}T(n) &= rT(n-1) + g(n) \\&= r(rT(n-2) + g(n-1)) + g(n) \\&= r^2T(n-2) + rg(n-1) + g(n) \\&= r^3T(n-3) + r^2g(n-2) + rg(n-1) + g(n) \\&\vdots \\&= r^nT(0) + \sum_{i=0}^{n-1} r^i g(n-i)\end{aligned}$$



First-Order Linear Recurrences

- **Theorem** For any positive constants a and r , and any function g defined on nonnegative integers, the solution to the first-order linear recurrence

$$T(n) = \begin{cases} rT(n-1) + g(n) & \text{if } n > 0 \\ a & \text{if } n = 0 \end{cases}$$

is

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i).$$



First-Order Linear Recurrences

- **Theorem** For any positive constants a and r , and any function g defined on nonnegative integers, the solution to the first-order linear recurrence

$$T(n) = \begin{cases} rT(n-1) + g(n) & \text{if } n > 0 \\ a & \text{if } n = 0 \end{cases}$$

is

$$T(n) = r^n a + \sum_{i=1}^n r^{n-i} g(i).$$

Proof by induction



Examples

- Solve $T(n) = 4T(n-1) + 2^n$ with $T(0) = 6$

Examples

- Solve $T(n) = 4T(n-1) + 2^n$ with $T(0) = 6$

$$\begin{aligned}T(n) &= 6 \cdot 4^n + \sum_{i=1}^n 4^{n-i} \cdot 2^i \\&= 6 \cdot 4^n + 4^n \sum_{i=1}^n 4^{-i} \cdot 2^i \\&= 6 \cdot 4^n + 4^n \sum_{i=1}^n \left(\frac{1}{2}\right)^i \\&= 6 \cdot 4^n + \left(1 - \frac{1}{2^n}\right) \cdot 4^n \\&= 7 \cdot 4^n - 2^n.\end{aligned}$$



Examples

- Solve $T(n) = 3T(n-1) + n$ with $T(0) = 10$



Examples

- Solve $T(n) = 3T(n-1) + n$ with $T(0) = 10$

$$\begin{aligned} T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\ &= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i} \end{aligned}$$

Examples

- Solve $T(n) = 3T(n-1) + n$ with $T(0) = 10$

$$\begin{aligned}T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\&= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i}\end{aligned}$$

Theorem. For any real number $x \neq 1$,

$$\sum_{i=1}^n ix^i = \frac{nx^{n+2} - (n+1)x^{n+1} + x}{(1-x)^2}.$$

Examples

- Solve $T(n) = 3T(n-1) + n$ with $T(0) = 10$

$$\begin{aligned}T(n) &= 10 \cdot 3^n + \sum_{i=1}^n 3^{n-i} \cdot i \\&= 10 \cdot 3^n + 3^n \sum_{i=1}^n i \cdot 3^{-i} \\&= 10 \cdot 3^n + 3^n \left(-\frac{3}{2}(n+1)3^{-(n+1)} - \frac{3}{4}3^{-(n+1)} + \frac{3}{4} \right) \\&= \frac{43}{4}3^n - \frac{n+1}{2} - \frac{1}{4}.\end{aligned}$$

Growth Rates of Solutions to Recurrences

- Divide and conquer algorithms
- Iterating recurrences
- Three different behaviors

Divide and conquer algorithms

- We just analyzed recurrences of the form

$$T(n) = \begin{cases} b & \text{if } n = 0 \\ r \cdot T(n-1) + a & \text{if } n > 0 \end{cases}$$



Divide and conquer algorithms

- We just analyzed recurrences of the form

$$T(n) = \begin{cases} b & \text{if } n = 0 \\ r \cdot T(n-1) + a & \text{if } n > 0 \end{cases}$$

These corresponded to the analysis of recursive algorithms in which a problem of size n is solved by recursively solving a problem of size $n - 1$.



Divide and conquer algorithms

- We just analyzed recurrences of the form

$$T(n) = \begin{cases} b & \text{if } n = 0 \\ r \cdot T(n-1) + a & \text{if } n > 0 \end{cases}$$

These corresponded to the analysis of recursive algorithms in which a problem of size n is solved by recursively solving a problem of size $n - 1$.

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

Divide and conquer algorithms

- We just analyzed recurrences of the form

$$T(n) = \begin{cases} b & \text{if } n = 0 \\ r \cdot T(n-1) + a & \text{if } n > 0 \end{cases}$$

These corresponded to the analysis of recursive algorithms in which a problem of size n is solved by recursively solving a problem of size $n-1$.

$$T(n) = r^n b + a \frac{1 - r^n}{1 - r}$$

We will now look at recurrences of the form

$$T(n) = \begin{cases} \text{something given} & \text{if } n \leq n_0 \\ r \cdot T(n/m) + a & \text{if } n > n_0 \end{cases}$$



Binary Search

- Someone has chosen a number x between 1 and n .
We need to discover x .



Binary Search

- Someone has chosen a number x between 1 and n . We need to discover x .

We are only allowed to ask **two types of questions**:

Binary Search

- Someone has chosen a number x between 1 and n . We need to discover x .

We are only allowed to ask two types of questions:

- ◇ Is x greater than k ?
- ◇ Is x equal to k ?

Binary Search

- Someone has chosen a number x between 1 and n . We need to discover x .

We are only allowed to ask two types of questions:

- ◇ Is x greater than k ?
- ◇ Is x equal to k ?

Our strategy will be to always ask **greater than** questions, at each step **halving our search range**, until the range only **contains one number**, when we ask a final **equal to** question.

Binary Search Example

1

32

48

64

33 - 1



Binary Search Example

1


32

48

64


Is $x > 32$?

Binary Search Example



Is $x > 32$? Answer: Yes


Binary Search Example



Is $x > 32$? Answer: Yes

Is $x > 48$?


Binary Search Example



Is $x > 32$? Answer: Yes

Is $x > 48$? Answer: No

Binary Search Example




Is $x > 32$? Answer: Yes

Is $x > 48$? Answer: No

Is $x > 40$?

Binary Search Example




Is $x > 32$? Answer: Yes

Is $x > 48$? Answer: No

Is $x > 40$? Answer: No

Binary Search Example




Is $x > 32$? Answer: Yes

Is $x > 48$? Answer: No

Is $x > 40$? Answer: No

Is $x > 36$?

Binary Search Example




Is $x > 32$? Answer: Yes

Is $x > 48$? Answer: No

Is $x > 40$? Answer: No

Is $x > 36$? Answer: No

Binary Search Example



Is $x > 32$? Answer: Yes


Is $x > 48$? Answer: No

Is $x > 40$? Answer: No

Is $x > 36$? Answer: No

Is $x > 34$?

Binary Search Example



Is $x > 32$? Answer: Yes


Is $x > 48$? Answer: No

Is $x > 40$? Answer: No

Is $x > 36$? Answer: No

Is $x > 34$? Answer: Yes

Binary Search Example



Is $x > 32$? Answer: Yes

Is $x > 48$? Answer: No


Is $x > 40$? Answer: No

Is $x > 36$? Answer: No

Is $x > 34$? Answer: Yes

Is $x > 35$?

Binary Search Example



Is $x > 32$? Answer: Yes

Is $x > 48$? Answer: No


Is $x > 40$? Answer: No

Is $x > 36$? Answer: No

Is $x > 34$? Answer: Yes

Is $x > 35$? Answer: No

Binary Search Example



Is $x > 32$? Answer: Yes

Is $x > 48$? Answer: No

Is $x > 40$? Answer: No


Is $x > 36$? Answer: No

Is $x > 34$? Answer: Yes

Is $x > 35$? Answer: No

Is $x = 35$?

Binary Search Example



Is $x > 32$?	Answer: Yes
Is $x > 48$?	Answer: No
Is $x > 40$?	Answer: No
Is $x > 36$?	Answer: No
Is $x > 34$?	Answer: Yes
Is $x > 35$?	Answer: No
Is $x = 35$?	Answer: BINGO!

Binary Search Example

- **Method**: Each guess **reduces** the problem to one in which the range is only **half** as big.



Binary Search Example

- **Method**: Each guess **reduces** the problem to one in which the range is only **half** as big.

This **divides** the original problem into one that is only half as big; we can now (**recursively**) **conquer** this smaller problem.



Binary Search Example

- **Method:** Each guess **reduces** the problem to one in which the range is only **half** as big.

This **divides** the original problem into one that is only half as big; we can now (**recursively**) **conquer** this smaller problem.

Note: When n is a power of 2, $T(n)$, the number of questions in a binary search on $[1, n]$, satisfies

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$



Binary Search Example

- **Method:** Each guess **reduces** the problem to one in which the range is only **half** as big.

This **divides** the original problem into one that is only half as big; we can now (**recursively**) **conquer** this smaller problem.

Note: When n is a power of 2, $T(n)$, the number of questions in a binary search on $[1, n]$, satisfies

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

This can also be proved **inductively**, similar to the tower of Hanoi recurrence.



Binary Search Example

- $T(n)$: number of questions in a binary search on $[1, n]$



Binary Search Example

- $T(n)$: number of questions in a binary search on $[1, n]$

Assume: n is a power of 2. Give recurrence for $T(n)$



Binary Search Example

- $T(n)$: number of questions in a binary search on $[1, n]$

Assume: n is a power of 2. Give recurrence for $T(n)$

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$



Binary Search Example

- $T(n)$: number of questions in a binary search on $[1, n]$

Assume: n is a power of 2. Give recurrence for $T(n)$

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

Number of questions needed for **binary search** on n items is:



Binary Search Example

- $T(n)$: number of questions in a binary search on $[1, n]$

Assume: n is a power of 2. Give recurrence for $T(n)$

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

Number of questions needed for **binary search** on n items is:

first step

+

time to perform binary search on the remaining $n/2$ items



Binary Search Example

- $T(n)$: number of questions in a binary search on $[1, n]$

Assume: n is a power of 2. Give recurrence for $T(n)$

$$T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

Number of questions needed for **binary search** on n items is:

first step

+

time to perform binary search on the remaining $n/2$ items

Base case (1 item): $T(1) = 1$ to ask: “**Is the number k ?**”



Binary Search Example

$$(*) \quad T(n) = \begin{cases} C_1 & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + C_2 & \text{if } n \geq 2 \end{cases}$$

For simplicity, we will (usually) assume that n is a power of 2 (or sometimes 3 or 4) and also often that constants such as C_1, C_2 are 1. This will let us replace a recurrence such as $(*)$ by one such as $(**)$.



Binary Search Example

$$(*) \quad T(n) = \begin{cases} C_1 & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + C_2 & \text{if } n \geq 2 \end{cases}$$

For simplicity, we will (usually) assume that n is a power of 2 (or sometimes 3 or 4) and also often that constants such as C_1, C_2 are 1. This will let us replace a recurrence such as (*) by one such as (**).

$$(**) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

Binary Search Example

$$(*) \quad T(n) = \begin{cases} C_1 & \text{if } n = 1 \\ T(\lceil n/2 \rceil) + C_2 & \text{if } n \geq 2 \end{cases}$$

For simplicity, we will (usually) assume that n is a power of 2 (or sometimes 3 or 4) and also often that constants such as C_1, C_2 are 1. This will let us replace a recurrence such as (*) by one such as (**).

$$(**) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

In practice, the solution of (*) will be very close to that of (**) (this can be proved mathematically). Hence, we can restrict attention to (**).

Growth Rates of Solutions to Recurrences

- Divide and conquer algorithms
- Iterating recurrences
- Three different behaviors

Iterating Recurrences: Example 1

■

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$



Iterating Recurrences: Example 1

■

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

This corresponds to solving a problem of size n , by

- (i) solving 2 subproblems of size $n/2$ and
- (ii) doing n units of additional work

or using $T(1)$ work for “bottom” case of $n = 1$



Iterating Recurrences: Example 1

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

This corresponds to solving a problem of size n , by

- (i) solving 2 subproblems of size $n/2$ and
- (ii) doing n units of additional work

or using $T(1)$ work for “bottom” case of $n = 1$

In the course “Analysis of Algorithms”, this is exactly how **Mergesort** works.



Iterating Recurrences: Example 1

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

This corresponds to solving a problem of size n , by

- (i) solving 2 subproblems of size $n/2$ and
- (ii) doing n units of additional work

or using $T(1)$ work for “bottom” case of $n = 1$

In the course “Analysis of Algorithms”, this is exactly how **Mergesort** works.

We now see how to solve $(*)$ by algebraically iterating the recurrence.



Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

$$T(n) = 2T\left(\frac{n}{2}\right) + n$$



Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that n is a power of 2

$$T(n) = 2T\left(\frac{n}{2}\right) + n = 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n$$



Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that n is a power of 2

$$\begin{aligned} T(n) &= 2T\left(\frac{n}{2}\right) + n &= 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n \\ &= 4T\left(\frac{n}{4}\right) + 2n &= 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n \end{aligned}$$



Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that n is a power of 2

$$\begin{aligned} T(n) &= 2T\left(\frac{n}{2}\right) + n &= 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n \\ &= 4T\left(\frac{n}{4}\right) + 2n &= 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n \\ &= 8T\left(\frac{n}{8}\right) + 3n \end{aligned}$$



Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that n is a power of 2

$$\begin{aligned}T(n) &= 2T\left(\frac{n}{2}\right) + n &= 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n \\&= 4T\left(\frac{n}{4}\right) + 2n &= 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n \\&= 8T\left(\frac{n}{8}\right) + 3n \\&\quad \vdots \quad \vdots \\&= 2^i T\left(\frac{n}{2^i}\right) + in\end{aligned}$$



Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that n is a power of 2

$$T(n) = 2T\left(\frac{n}{2}\right) + n = 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n$$

$$= 4T\left(\frac{n}{4}\right) + 2n = 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n$$

$$= 8T\left(\frac{n}{8}\right) + 3n$$

$$\vdots \quad \vdots$$
$$= 2^i T\left(\frac{n}{2^i}\right) + in$$

$$\vdots \quad \vdots$$
$$= 2^{\log_2 n} T\left(\frac{n}{2^{\log_2 n}}\right) + (\log_2 n)n$$

End when $i = \log_2 n$



Iterating Recurrences: Example 1

- Algebraically iterating the recurrence

Assume that n is a power of 2

$$T(n) = 2T\left(\frac{n}{2}\right) + n = 2\left(2T\left(\frac{n}{4}\right) + \frac{n}{2}\right) + n$$

$$= 4T\left(\frac{n}{4}\right) + 2n = 4\left(2T\left(\frac{n}{8}\right) + \frac{n}{4}\right) + 2n$$

$$= 8T\left(\frac{n}{8}\right) + 3n$$

$$\vdots \quad \vdots$$
$$= 2^i T\left(\frac{n}{2^i}\right) + in$$

End when $i = \log_2 n$

$$\vdots \quad \vdots$$
$$= 2^{\log_2 n} T\left(\frac{n}{2^{\log_2 n}}\right) + (\log_2 n)n$$

$$= nT(1) + n\log_2 n$$

Iterating Recurrences: Example 1

- We just iterated the recurrence to derive that the solution to

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

is $nT(1) + n \log_2 n$.



Iterating Recurrences: Example 1

- We just iterated the recurrence to derive that the solution to

$$(*) \quad T(n) = \begin{cases} T(1) & \text{if } n = 1 \\ 2T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

is $nT(1) + n \log_2 n$.

Note: Technically, we still need to use **induction** to prove that our solution is correct. Practically, we **never** explicitly perform this step, since it is obvious how the induction would work.



Iterating Recurrences: Example 2

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$



Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1$$



Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1 = \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1$$



Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + 1 &&= \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1 \\ &= T\left(\frac{n}{2^2}\right) + 2 \end{aligned}$$



Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + 1 &= \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1 \\ &= T\left(\frac{n}{2^2}\right) + 2 &= \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2 \end{aligned}$$



Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + 1 &&= \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1 \\ &= T\left(\frac{n}{2^2}\right) + 2 &&= \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2 \\ &= T\left(\frac{n}{2^3}\right) + 3 \end{aligned}$$



Iterating Recurrences: Example 2



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + 1 &&= \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1 \\ &= T\left(\frac{n}{2^2}\right) + 2 &&= \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2 \\ &= T\left(\frac{n}{2^3}\right) + 3 \\ &\quad \vdots \quad \quad \quad \vdots \\ &= T\left(\frac{n}{2^i}\right) + i \end{aligned}$$



Iterating Recurrences: Example 2

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1 = \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1$$

$$= T\left(\frac{n}{2^2}\right) + 2 = \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2$$

$$= T\left(\frac{n}{2^3}\right) + 3$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^i}\right) + i \end{array}$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^{\log_2 n}}\right) + \log_2 n \end{array}$$



Iterating Recurrences: Example 2

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + 1 & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + 1 = \left(T\left(\frac{n}{2^2}\right) + 1\right) + 1$$

$$= T\left(\frac{n}{2^2}\right) + 2 = \left(T\left(\frac{n}{2^3}\right) + 1\right) + 2$$

$$= T\left(\frac{n}{2^3}\right) + 3$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^i}\right) + i \end{array}$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^{\log_2 n}}\right) + \log_2 n = 1 + \log_2 n \end{array}$$



Iterating Recurrences: Example 3

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$



Iterating Recurrences: Example 3



$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + n \\ &= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n \end{aligned}$$



Iterating Recurrences: Example 3

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + n \\ &= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n \\ &= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n \end{aligned}$$



Iterating Recurrences: Example 3

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= T\left(\frac{n}{2}\right) + n \\ &= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n \\ &= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n \\ &\quad \vdots \\ &= T\left(\frac{n}{2^i}\right) + \frac{n}{2^{i-1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n \end{aligned}$$



Iterating Recurrences: Example 3

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + n$$

$$= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n$$

$$= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^i}\right) + \frac{n}{2^{i-1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n \end{array}$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{n}{2^{\log_2 n - 1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n \end{array}$$



Iterating Recurrences: Example 3

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + n$$

$$= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n$$

$$= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^i}\right) + \frac{n}{2^{i-1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n \end{array}$$

$$\begin{array}{c} \vdots \\ \vdots \\ = T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{n}{2^{\log_2 n - 1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n \end{array}$$

$$= 1 + 2 + 2^2 + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$

Iterating Recurrences: Example 3

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = T\left(\frac{n}{2}\right) + n$$

$$= T\left(\frac{n}{2^2}\right) + \frac{n}{2} + n$$

$$= T\left(\frac{n}{2^3}\right) + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^i}\right) + \frac{n}{2^{i-1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$

$$\vdots \quad \vdots$$

$$= T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{n}{2^{\log_2 n - 1}} + \cdots + \frac{n}{2^2} + \frac{n}{2} + n$$

$$= 1 + 2 + 2^2 + \cdots + \frac{n}{2^2} + \frac{n}{2} + n = \Theta(n)$$

Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$



Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$T(n) = 3T\left(\frac{n}{3}\right) + n$$



Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$T(n) = 3T\left(\frac{n}{3}\right) + n = 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n$$



Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n \end{aligned}$$



Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \end{aligned}$$



Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \\ &= 3^3 T\left(\frac{n}{3^3}\right) + 3n \end{aligned}$$



Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \\ &= 3^3 T\left(\frac{n}{3^3}\right) + 3n \\ &\quad \vdots \quad \quad \quad \vdots \\ &= 3^i T\left(\frac{n}{3^i}\right) + in \end{aligned}$$



Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \\ &= 3^3 T\left(\frac{n}{3^3}\right) + 3n \\ &\quad \vdots \quad \vdots \\ &= 3^i T\left(\frac{n}{3^i}\right) + in \\ &\quad \vdots \quad \vdots \\ &= 3^{\log_3 n} T\left(\frac{n}{3^{\log_3 n}}\right) + n \log_3 n \end{aligned}$$



Iterating Recurrences: Example 4

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n < 3 \\ 3T(n/3) + n & \text{if } n \geq 3 \end{cases}$$

$$\begin{aligned} T(n) &= 3T\left(\frac{n}{3}\right) + n &= 3\left(3T\left(\frac{n}{3^2}\right) + \frac{n}{3}\right) + n \\ &= 3^2 T\left(\frac{n}{3^2}\right) + 2n &= 3^2\left(3T\left(\frac{n}{3^3}\right) + \frac{n}{3^2}\right) + 2n \\ &= 3^3 T\left(\frac{n}{3^3}\right) + 3n \\ &\quad \vdots \quad \vdots \\ &= 3^i T\left(\frac{n}{3^i}\right) + in \\ &\quad \vdots \quad \vdots \\ &= 3^{\log_3 n} T\left(\frac{n}{3^{\log_3 n}}\right) + n \log_3 n = n + n \log_3 n \end{aligned}$$



Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$



Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = 4T\left(\frac{n}{2}\right) + n$$



Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$T(n) = 4T\left(\frac{n}{2}\right) + n = 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n$$



Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n \end{aligned}$$



Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \end{aligned}$$



Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &&= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &&= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \\ &= 4^3 T\left(\frac{n}{2^3}\right) + \frac{4^2}{2^2}n + \frac{4}{2}n + n \end{aligned}$$



Iterating Recurrences: Example 5

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &&= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &&= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \\ &= 4^3 T\left(\frac{n}{2^3}\right) + \frac{4^2}{2^2}n + \frac{4}{2}n + n \\ &\quad \vdots \quad \quad \quad \vdots \\ &= 4^i T\left(\frac{n}{2^i}\right) + \frac{4^{i-1}}{2^{i-1}}n + \cdots + \frac{4^2}{2^2}n + n \end{aligned}$$

Iterating Recurrences: Example 5

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &&= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &&= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \\ &= 4^3T\left(\frac{n}{2^3}\right) + \frac{4^2}{2^2}n + \frac{4}{2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^iT\left(\frac{n}{2^i}\right) + \frac{4^{i-1}}{2^{i-1}}n + \cdots + \frac{4^2}{2^2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^{\log_2 n}T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{4^{\log_2 n - 1}}{2^{\log_2 n - 1}}n + \cdots + \frac{4}{2}n + n \end{aligned}$$

Iterating Recurrences: Example 5

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$$\begin{aligned} T(n) &= 4T\left(\frac{n}{2}\right) + n &&= 4\left(4T\left(\frac{n}{2^2}\right) + \frac{n}{2}\right) + n \\ &= 4^2 T\left(\frac{n}{2^2}\right) + \frac{4}{2}n + n &&= 4^2\left(4T\left(\frac{n}{2^3}\right) + \frac{n}{2^2}\right) + \frac{4}{2}n + n \\ &= 4^3 T\left(\frac{n}{2^3}\right) + \frac{4^2}{2^2}n + \frac{4}{2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^i T\left(\frac{n}{2^i}\right) + \frac{4^{i-1}}{2^{i-1}}n + \cdots + \frac{4^2}{2^2}n + n \\ &\quad \vdots \quad \vdots \\ &= 4^{\log_2 n} T\left(\frac{n}{2^{\log_2 n}}\right) + \frac{4^{\log_2 n - 1}}{2^{\log_2 n - 1}}n + \cdots + \frac{4}{2}n + n \\ &= 2n^2 - n \end{aligned}$$



Three Different Behaviors

- Compare the iteration for the recurrences

$$T(n) = 2T(n/2) + n$$

$$T(n) = T(n/2) + n$$

$$T(n) = 4T(n/2) + n$$



Three Different Behaviors

- Compare the iteration for the recurrences

$$T(n) = 2T(n/2) + n$$

$$T(n) = T(n/2) + n$$

$$T(n) = 4T(n/2) + n$$

- ◇ all three recurrences iterate $\log_2 n$ times
- ◇ in each case, size of subproblem in next iteration is **half** the size in the preceding iteration level



Three Different Behaviors

- **Theorem** Suppose that we have a recurrence of the form

$$T(n) = aT(n/2) + n,$$

where a is a positive integer and $T(1)$ is nonnegative. Then we have the following **big Θ** bounds on the solution:

1. If $a < 2$, then $T(n) = \Theta(n)$.
2. If $a = 2$, then $T(n) = \Theta(n \log n)$.
3. If $a > 2$, then $T(n) = \Theta(n^{\log_2 a})$



Three Different Behaviors

- **Theorem** Suppose that we have a recurrence of the form

$$T(n) = aT(n/2) + n,$$

where a is a positive integer and $T(1)$ is nonnegative. Then we have the following **big Θ** bounds on the solution:

1. If $a < 2$, then $T(n) = \Theta(n)$.
2. If $a = 2$, then $T(n) = \Theta(n \log n)$.
3. If $a > 2$, then $T(n) = \Theta(n^{\log_2 a})$

Proof

We already proved Case 1 when $a = 1$ in Example 3.

(will not prove it for $1 < a < 2$)

We already proved Case 2 in Example 1.

We will now prove Case 3.



Iterating Recurrences

- $T(n) = aT(n/2) + n$, where $a > 2$. Assume that $n = 2^i$.



Iterating Recurrences

- $T(n) = aT(n/2) + n$, where $a > 2$. Assume that $n = 2^i$.

Iterating as in Example 5 gives

$$T(n) = a^i T\left(\frac{n}{2^i}\right) + \left(\frac{a^{i-1}}{2^{i-1}} + \frac{a^{i-2}}{2^{i-2}} + \cdots + \frac{a}{2} + 1\right) n$$



Iterating Recurrences

- $T(n) = aT(n/2) + n$, where $a > 2$. Assume that $n = 2^i$.

Iterating as in Example 5 gives

$$T(n) = a^i T\left(\frac{n}{2^i}\right) + \left(\frac{a^{i-1}}{2^{i-1}} + \frac{a^{i-2}}{2^{i-2}} + \cdots + \frac{a}{2} + 1\right) n$$

$$T(n) = a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$

Work at
“bottom”

Iterated
Work



Total work

- The total work is

$$T(n) = a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$



Total work

- The total work is

$$T(n) = a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$

Since $a > 2$, the geometric series is Θ of the largest term.

$$n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i = n \frac{1 - (a/2)^{\log_2 n}}{1 - a/2} = n\Theta((a/2)^{\log_2 n - 1})$$



Total work

- n times the largest term in the geometric series is

$$n \left(\frac{a}{2}\right)^{\log_2 n - 1} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{2^{\log_2 n}} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{n} = \frac{2}{a} \cdot a^{\log_2 n}$$

Total work

- n times the largest term in the geometric series is

$$n \left(\frac{a}{2}\right)^{\log_2 n - 1} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{2^{\log_2 n}} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{n} = \frac{2}{a} \cdot a^{\log_2 n}$$

Notice that

$$a^{\log_2 n} = (2^{\log_2 a})^{\log_2 n} = (2^{\log_2 n})^{\log_2 a} = n^{\log_2 a}$$

Total work

- n times the largest term in the geometric series is

$$n \left(\frac{a}{2}\right)^{\log_2 n - 1} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{2^{\log_2 n}} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{n} = \frac{2}{a} \cdot a^{\log_2 n}$$

Notice that

$$a^{\log_2 n} = (2^{\log_2 a})^{\log_2 n} = (2^{\log_2 n})^{\log_2 a} = n^{\log_2 a}$$

So the total work is

$$a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$

Total work

- n times the largest term in the geometric series is

$$n \left(\frac{a}{2}\right)^{\log_2 n - 1} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{2^{\log_2 n}} = \frac{2}{a} \cdot \frac{n \cdot a^{\log_2 n}}{n} = \frac{2}{a} \cdot a^{\log_2 n}$$

Notice that

$$a^{\log_2 n} = (2^{\log_2 a})^{\log_2 n} = (2^{\log_2 n})^{\log_2 a} = n^{\log_2 a}$$

So the total work is

$$a^{\log_2 n} T(1) + n \sum_{i=0}^{\log_2 n - 1} \left(\frac{a}{2}\right)^i$$

$$\Theta(n^{\log_2 a})$$

$$\Theta(n^{\log_2 a})$$



Example 5 Recap

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$



Example 5 Recap

■

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$a = 4$, so the Theorem says that

$$T(n) = \Theta(n^{\log_2 a}) = \Theta(n^{\log_2 4}) = \Theta(n^2)$$



Example 5 Recap

$$(*) \quad T(n) = \begin{cases} 1 & \text{if } n = 1 \\ 4T(n/2) + n & \text{if } n \geq 2 \end{cases}$$

$a = 4$, so the Theorem says that

$$T(n) = \Theta(n^{\log_2 a}) = \Theta(n^{\log_2 4}) = \Theta(n^2)$$

This matches with the exact answer of $2n^2 - n$.

Three Different Behaviors

- **Theorem** Suppose that we have a recurrence of the form

$$T(n) = aT(n/2) + n,$$

where a is a positive integer and $T(1)$ is nonnegative. Then we have the following big Θ bounds on the solution:

1. If $a < 2$, then $T(n) = \Theta(n)$.
2. If $a = 2$, then $T(n) = \Theta(n \log n)$.
3. If $a > 2$, then $T(n) = \Theta(n^{\log_2 a})$.



The Master Theorem

- **Theorem** Suppose that we have a recurrence of the form

$$T(n) = aT(n/b) + cn^d,$$

where a is a positive integer, $b \geq 1$, c, d are real numbers with c positive and d nonnegative, and $T(1)$ is nonnegative. Then we have the following **big Θ** bounds on the solution:

1. If $a < b^d$, then $T(n) = \Theta(n^d)$.
2. If $a = b^d$, then $T(n) = \Theta(n^d \log n)$.
3. If $a > b^d$, then $T(n) = \Theta(n^{\log_b a})$



Next Lecture

- counting ...

