



# CS215 DISCRETE MATH

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: [wangqi@sustech.edu.cn](mailto:wangqi@sustech.edu.cn)

# Binary Relation

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .



# Binary Relation

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .

- **Definition:** A *relation on the set  $A$*  is a relation **from  $A$  to itself**.

# Binary Relation

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .

- **Definition:** A *relation on the set  $A$*  is a relation **from  $A$  to itself**.
- **Theorem** The number of binary relations on a set  $A$ , where  $|A| = n$  is  $2^{n^2}$ .



# Properties of Relations

- **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .



# Properties of Relations

- **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .
- **Irreflexive Relation:** A relation  $R$  on a set  $A$  is called *irreflexive* if  $(a, a) \notin R$  for **every** element  $a \in A$ .

# Properties of Relations

- **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .
- **Irreflexive Relation:** A relation  $R$  on a set  $A$  is called *irreflexive* if  $(a, a) \notin R$  for **every** element  $a \in A$ .
- **Symmetric Relation:** A relation  $R$  on a set  $A$  is called *symmetric* if  $(b, a) \in R$  whenever  $(a, b) \in R$  for **all**  $a, b \in A$ .



# Properties of Relations

- **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .
- **Irreflexive Relation:** A relation  $R$  on a set  $A$  is called *irreflexive* if  $(a, a) \notin R$  for **every** element  $a \in A$ .
- **Symmetric Relation:** A relation  $R$  on a set  $A$  is called *symmetric* if  $(b, a) \in R$  whenever  $(a, b) \in R$  for **all**  $a, b \in A$ .
- **Antisymmetric Relation:** A relation  $R$  on a set  $A$  is called *antisymmetric* if  $(b, a) \in R$  and  $(a, b) \in R$  implies  $a = b$  for **all**  $a, b \in A$ .



# Properties of Relations

- **Transitive Relation:** A relation  $R$  on a set  $A$  is called *transitive* if  $(a, b) \in R$  and  $(b, c) \in R$  implies  $(a, c) \in R$  for **all**  $a, b, c \in A$ .



# Properties of Relations

- **Transitive Relation:** A relation  $R$  on a set  $A$  is called *transitive* if  $(a, b) \in R$  and  $(b, c) \in R$  implies  $(a, c) \in R$  for **all**  $a, b, c \in A$ .

**Example:** Assume that  $R_{div} = \{(a, b) : a|b\}$  on  $A = \{1, 2, 3, 4\}$ .



# Properties of Relations

- **Transitive Relation:** A relation  $R$  on a set  $A$  is called *transitive* if  $(a, b) \in R$  and  $(b, c) \in R$  implies  $(a, c) \in R$  for **all**  $a, b, c \in A$ .

**Example:** Assume that  $R_{div} = \{(a, b) : a|b\}$  on  $A = \{1, 2, 3, 4\}$ .

Is  $R_{div}$  *transitive*?



# Properties of Relations

- **Transitive Relation:** A relation  $R$  on a set  $A$  is called *transitive* if  $(a, b) \in R$  and  $(b, c) \in R$  implies  $(a, c) \in R$  for **all**  $a, b, c \in A$ .

**Example:** Assume that  $R_{div} = \{(a, b) : a|b\}$  on  $A = \{1, 2, 3, 4\}$ .

Is  $R_{div}$  *transitive*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$



# Properties of Relations

- **Transitive Relation:** A relation  $R$  on a set  $A$  is called *transitive* if  $(a, b) \in R$  and  $(b, c) \in R$  implies  $(a, c) \in R$  for **all**  $a, b, c \in A$ .

**Example:** Assume that  $R_{div} = \{(a, b) : a|b\}$  on  $A = \{1, 2, 3, 4\}$ .

Is  $R_{div}$  *transitive*?

$$R_{div} = \{(1, 1), (1, 2), (1, 3), (1, 4), (2, 2), (2, 4), (3, 3), (4, 4)\}.$$

**Yes.** If  $a|b$  and  $b|c$ , then  $a|c$ .



# Transitive Relation

- **Example:** Assume that  $R_{\neq} = \{(a, b) : a \neq b\}$  on  $A = \{1, 2, 3, 4\}$ .

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

# Transitive Relation

- **Example:** Assume that  $R_{\neq} = \{(a, b) : a \neq b\}$  on  $A = \{1, 2, 3, 4\}$ .

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is  $R_{\neq}$  transitive?



# Transitive Relation

- **Example:** Assume that  $R_{\neq} = \{(a, b) : a \neq b\}$  on  $A = \{1, 2, 3, 4\}$ .

$$R_{\neq} = \{(1, 2), (1, 3), (1, 4), (2, 1), (2, 3), (2, 4), \\ (3, 1), (3, 2), (3, 4), (4, 1), (4, 2), (4, 3)\}.$$

Is  $R_{\neq}$  transitive?

**No.**  $(1, 2), (2, 1) \in R_{\neq}$  but  $(1, 1) \notin R_{\neq}$ .





# Transitive Relation

- **Example:** Assume that  $R = \{(1, 2), (2, 2), (3, 3)\}$  on  $A = \{1, 2, 3, 4\}$ .

# Transitive Relation

- **Example:** Assume that  $R = \{(1, 2), (2, 2), (3, 3)\}$  on  $A = \{1, 2, 3, 4\}$ .

Is  $R$  transitive?

# Transitive Relation

- **Example:** Assume that  $R = \{(1, 2), (2, 2), (3, 3)\}$  on  $A = \{1, 2, 3, 4\}$ .

Is  $R$  transitive?

Yes.

# Combining Relations

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .



# Combining Relations

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .

**Combining Relations:** Since **relations are sets**, we can *combine* relations via **set operations**.



# Combining Relations

- **Definition:** Let  $A$  and  $B$  be two sets. A *binary relation from  $A$  to  $B$*  is a **subset** of a **Cartesian product**  $A \times B$ .

Let  $R \subseteq A \times B$  denote  $R$  is a set of ordered pairs of the form  $(a, b)$  where  $a \in A$  and  $b \in B$ .

**Combining Relations:** Since **relations are sets**, we can *combine* relations via **set operations**.

Set operations: **union, intersection, difference, etc.**



# Combining Relations

- **Example:** Let  $A = \{1, 2, 3\}$ ,  $B = \{u, v\}$ , and  
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$ ,  
 $R_2 = \{(1, v), (3, u), (3, v)\}$

# Combining Relations

- **Example:** Let  $A = \{1, 2, 3\}$ ,  $B = \{u, v\}$ , and  
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$ ,  
 $R_2 = \{(1, v), (3, u), (3, v)\}$

What is  $R_1 \cup R_2$ ,  $R_1 \cap R_2$ ,  $R_1 - R_2$ ,  $R_2 - R_1$ ?





# Combining Relations

- **Example:** Let  $A = \{1, 2, 3\}$ ,  $B = \{u, v\}$ , and  
 $R_1 = \{(1, u), (2, u), (2, v), (3, u)\}$ ,  
 $R_2 = \{(1, v), (3, u), (3, v)\}$

What is  $R_1 \cup R_2$ ,  $R_1 \cap R_2$ ,  $R_1 - R_2$ ,  $R_2 - R_1$ ?

We may also combine relations by **matrix operations**.

# Composite of Relations

- **Definition:** Let  $R$  be a relation from a set  $A$  to a set  $B$  and  $S$  be a relation from  $B$  to  $C$ . The *composite of  $R$  and  $S$*  is the relation consisting of the ordered pairs  $(a, c)$  where  $a \in A$  and  $c \in C$  and for which there is a  $b \in B$  such that  $(a, b) \in R$  and  $(b, c) \in S$ . We denote the composite of  $R$  and  $S$  by  $S \circ R$ .



# Composite of Relations

- **Definition:** Let  $R$  be a relation from a set  $A$  to a set  $B$  and  $S$  be a relation from  $B$  to  $C$ . The *composite of  $R$  and  $S$*  is the relation consisting of the ordered pairs  $(a, c)$  where  $a \in A$  and  $c \in C$  and for which there is a  $b \in B$  such that  $(a, b) \in R$  and  $(b, c) \in S$ . We denote the composite of  $R$  and  $S$  by  $S \circ R$ .

**Example:** Let  $A = \{1, 2, 3\}$ ,  $B = \{0, 1, 2\}$ , and  $C = \{a, b\}$



# Composite of Relations

- **Definition:** Let  $R$  be a relation from a set  $A$  to a set  $B$  and  $S$  be a relation from  $B$  to  $C$ . The *composite of  $R$  and  $S$*  is the relation consisting of the ordered pairs  $(a, c)$  where  $a \in A$  and  $c \in C$  and for which there is a  $b \in B$  such that  $(a, b) \in R$  and  $(b, c) \in S$ . We denote the composite of  $R$  and  $S$  by  $S \circ R$ .

**Example:** Let  $A = \{1, 2, 3\}$ ,  $B = \{0, 1, 2\}$ , and  $C = \{a, b\}$

$$R = \{(1, 0), (1, 2), (3, 1), (3, 2)\}$$

$$S = \{(0, b), (1, a), (2, b)\}$$



# Composite of Relations

- **Definition:** Let  $R$  be a relation from a set  $A$  to a set  $B$  and  $S$  be a relation from  $B$  to  $C$ . The *composite of  $R$  and  $S$*  is the relation consisting of the ordered pairs  $(a, c)$  where  $a \in A$  and  $c \in C$  and for which there is a  $b \in B$  such that  $(a, b) \in R$  and  $(b, c) \in S$ . We denote the composite of  $R$  and  $S$  by  $S \circ R$ .

**Example:** Let  $A = \{1, 2, 3\}$ ,  $B = \{0, 1, 2\}$ , and  $C = \{a, b\}$

$$R = \{(1, 0), (1, 2), (3, 1), (3, 2)\}$$

$$S = \{(0, b), (1, a), (2, b)\}$$

$$S \circ R = \{(1, b), (3, a), (3, b)\}$$



# Implementation of Composite

- **Example:** Let  $A = \{1, 2\}$ ,  $B = \{1, 2, 3\}$ , and  $C = \{a, b\}$   
 $R = \{(1, 2), (1, 3), (2, 1)\}$  is a relation from  $A$  to  $B$   
 $S = \{(1, a), (3, b), (3, a)\}$  is a relation from  $B$  to  $C$



# Implementation of Composite

- **Example:** Let  $A = \{1, 2\}$ ,  $B = \{1, 2, 3\}$ , and  $C = \{a, b\}$   
 $R = \{(1, 2), (1, 3), (2, 1)\}$  is a relation from  $A$  to  $B$   
 $S = \{(1, a), (3, b), (3, a)\}$  is a relation from  $B$  to  $C$   
 $S \circ R = \{(1, b), (1, a), (2, a)\}$



# Implementation of Composite

■ **Example:** Let  $A = \{1, 2\}$ ,  $B = \{1, 2, 3\}$ , and  $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$  is a relation from  $A$  to  $B$

$S = \{(1, a), (3, b), (3, a)\}$  is a relation from  $B$  to  $C$

$$S \circ R = \{(1, b), (1, a), (2, a)\}$$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix}$$



# Implementation of Composite

■ **Example:** Let  $A = \{1, 2\}$ ,  $B = \{1, 2, 3\}$ , and  $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$  is a relation from  $A$  to  $B$

$S = \{(1, a), (3, b), (3, a)\}$  is a relation from  $B$  to  $C$

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 \\ 3 \end{matrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix}$$



# Implementation of Composite

■ **Example:** Let  $A = \{1, 2\}$ ,  $B = \{1, 2, 3\}$ , and  $C = \{a, b\}$

$R = \{(1, 2), (1, 3), (2, 1)\}$  is a relation from  $A$  to  $B$

$S = \{(1, a), (3, b), (3, a)\}$  is a relation from  $B$  to  $C$

$S \circ R = \{(1, b), (1, a), (2, a)\}$

$$M_R = \begin{matrix} & \begin{matrix} 1 & 2 & 3 \end{matrix} \\ \begin{matrix} 1 \\ 2 \end{matrix} & \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix} \end{matrix} \quad M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 & 3 \end{matrix} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{matrix}$$

$$M_R \odot M_S = \begin{matrix} & \begin{matrix} a & b \end{matrix} \\ \begin{matrix} 1 & 2 \end{matrix} & \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \end{matrix}$$



# Composite of Relations

- **Definition** Let  $R$  be a relation on  $A$ . The *powers*  $R^n$ , for  $n = 1, 2, 3, \dots$ , is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$



# Composite of Relations

- **Definition** Let  $R$  be a relation on  $A$ . The *powers*  $R^n$ , for  $n = 1, 2, 3, \dots$ , is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

**Example:** Let  $A = \{1, 2, 3, 4\}$ , and  
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$



# Composite of Relations

- **Definition** Let  $R$  be a relation on  $A$ . The *powers*  $R^n$ , for  $n = 1, 2, 3, \dots$ , is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

**Example:** Let  $A = \{1, 2, 3, 4\}$ , and  
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$



# Composite of Relations

- **Definition** Let  $R$  be a relation on  $A$ . The *powers*  $R^n$ , for  $n = 1, 2, 3, \dots$ , is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

**Example:** Let  $A = \{1, 2, 3, 4\}$ , and  
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$



# Composite of Relations

- **Definition** Let  $R$  be a relation on  $A$ . The *powers*  $R^n$ , for  $n = 1, 2, 3, \dots$ , is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

**Example:** Let  $A = \{1, 2, 3, 4\}$ , and  
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$



# Composite of Relations

- **Definition** Let  $R$  be a relation on  $A$ . The *powers*  $R^n$ , for  $n = 1, 2, 3, \dots$ , is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

**Example:** Let  $A = \{1, 2, 3, 4\}$ , and  
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^4 = R^3 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$





# Composite of Relations

- **Definition** Let  $R$  be a relation on  $A$ . The *powers*  $R^n$ , for  $n = 1, 2, 3, \dots$ , is defined **inductively** by

$$R^1 = R \text{ and } R^{n+1} = R^n \circ R$$

**Example:** Let  $A = \{1, 2, 3, 4\}$ , and  
 $R = \{(1, 2), (2, 3), (2, 4), (3, 3)\}$

$$R^1 = R$$

$$R^2 = R \circ R = \{(1, 3), (1, 4), (2, 3), (3, 3)\}$$

$$R^3 = R^2 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^4 = R^3 \circ R = \{(1, 3), (2, 3), (3, 3)\}$$

$$R^k = ? \text{ for } k > 3$$



# Transitive Relation and $R^n$

- **Theorem** The relation  $R$  on a set  $A$  is transitive if and only if  $R^n \subseteq R$  for  $n = 1, 2, 3, \dots$



# Transitive Relation and $R^n$

- **Theorem** The relation  $R$  on a set  $A$  is transitive if and only if  $R^n \subseteq R$  for  $n = 1, 2, 3, \dots$

**Proof.**



# Transitive Relation and $R^n$

- **Theorem** The relation  $R$  on a set  $A$  is transitive if and only if  $R^n \subseteq R$  for  $n = 1, 2, 3, \dots$

**Proof.**

“if” part: In particular,  $R^2 \subseteq R$ .



# Transitive Relation and $R^n$

- **Theorem** The relation  $R$  on a set  $A$  is transitive if and only if  $R^n \subseteq R$  for  $n = 1, 2, 3, \dots$

## Proof.

“if” part: In particular,  $R^2 \subseteq R$ .

If  $(a, b) \in R$  and  $(b, c) \in R$ , then by the definition of composition, we have  $(a, c) \in R^2 \subseteq R$ .



# Transitive Relation and $R^n$

- **Theorem** The relation  $R$  on a set  $A$  is transitive if and only if  $R^n \subseteq R$  for  $n = 1, 2, 3, \dots$

## Proof.

“if” part: In particular,  $R^2 \subseteq R$ .

If  $(a, b) \in R$  and  $(b, c) \in R$ , then by the definition of composition, we have  $(a, c) \in R^2 \subseteq R$ .

“only if” part: by induction.



# Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set  $A$  with  $|A| = n$  is  $2^{n(n-1)}$ .



# Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set  $A$  with  $|A| = n$  is  $2^{n(n-1)}$ .

**Proof.** A reflexive relation  $R$  on  $A$  must contain all pairs  $(a, a)$  for every  $a \in A$ .



# Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set  $A$  with  $|A| = n$  is  $2^{n(n-1)}$ .

**Proof.** A reflexive relation  $R$  on  $A$  must contain all pairs  $(a, a)$  for every  $a \in A$ .

All other pairs in  $R$  are of the form  $(a, b)$  with  $a \neq b$  s.t.  $a, b \in A$ .



# Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set  $A$  with  $|A| = n$  is  $2^{n(n-1)}$ .

**Proof.** A reflexive relation  $R$  on  $A$  must contain all pairs  $(a, a)$  for every  $a \in A$ .

All other pairs in  $R$  are of the form  $(a, b)$  with  $a \neq b$  s.t.  $a, b \in A$ .

How many of these pairs are there?



# Number of Reflexive Relations

- **Theorem** The number of reflexive relations on a set  $A$  with  $|A| = n$  is  $2^{n(n-1)}$ .

**Proof.** A reflexive relation  $R$  on  $A$  must contain all pairs  $(a, a)$  for every  $a \in A$ .

All other pairs in  $R$  are of the form  $(a, b)$  with  $a \neq b$  s.t.  $a, b \in A$ .

How many of these pairs are there?

How many subsets on  $n(n-1)$  elements are there?



# Summary on Properties of Relations

- **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .



# Summary on Properties of Relations

- **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .

**Irreflexive Relation:** A relation  $R$  on a set  $A$  is called *irreflexive* if  $(a, a) \notin R$  for **every** element  $a \in A$ .



# Summary on Properties of Relations

■ **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .

**Irreflexive Relation:** A relation  $R$  on a set  $A$  is called *irreflexive* if  $(a, a) \notin R$  for **every** element  $a \in A$ .

**Symmetric Relation:** A relation  $R$  on a set  $A$  is called *symmetric* if  $(b, a) \in R$  whenever  $(a, b) \in R$  for **all**  $a, b \in A$ .

# Summary on Properties of Relations

- **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .

**Irreflexive Relation:** A relation  $R$  on a set  $A$  is called *irreflexive* if  $(a, a) \notin R$  for **every** element  $a \in A$ .

**Symmetric Relation:** A relation  $R$  on a set  $A$  is called *symmetric* if  $(b, a) \in R$  whenever  $(a, b) \in R$  for **all**  $a, b \in A$ .

**Antisymmetric Relation:** A relation  $R$  on a set  $A$  is called *antisymmetric* if  $(b, a) \in R$  and  $(a, b) \in R$  implies  $a = b$  for **all**  $a, b \in A$ .



# Summary on Properties of Relations

■ **Reflexive Relation:** A relation  $R$  on a set  $A$  is called *reflexive* if  $(a, a) \in R$  for **every** element  $a \in A$ .

**Irreflexive Relation:** A relation  $R$  on a set  $A$  is called *irreflexive* if  $(a, a) \notin R$  for **every** element  $a \in A$ .

**Symmetric Relation:** A relation  $R$  on a set  $A$  is called *symmetric* if  $(b, a) \in R$  whenever  $(a, b) \in R$  for **all**  $a, b \in A$ .

**Antisymmetric Relation:** A relation  $R$  on a set  $A$  is called *antisymmetric* if  $(b, a) \in R$  and  $(a, b) \in R$  implies  $a = b$  for **all**  $a, b \in A$ .

**Transitive Relation:** A relation  $R$  on a set  $A$  is called *transitive* if  $(a, b) \in R$  and  $(b, c) \in R$  implies  $(a, c) \in R$  for **all**  $a, b, c \in A$ .



# Representing Relations

- Some ways to represent *n-ary* relations:
  - with an *explicit list* or *table* of its tuples
  - with a *function* from the domain to  $\{T, F\}$


# Representing Relations

- Some ways to represent *n-ary* relations:
  - with an *explicit list* or *table* of its tuples
  - with a *function* from the domain to  $\{T, F\}$
- Some special ways to represent *binary* relations:
  - with a *zero-one matrix*
  - with a *directed graph*




# Representing Relations

- Some ways to represent *n-ary* relations:
  - with an *explicit list* or *table* of its tuples
  - with a *function* from the domain to  $\{T, F\}$
- Some special ways to represent *binary* relations:
  - with a *zero-one matrix*
  - with a *directed graph*




# Representing Relations


- Some ways to represent *n-ary* relations:
  - with an *explicit list* or *table* of its tuples
  - with a *function* from the domain to  $\{T, F\}$
- Some special ways to represent *binary* relations:
  - with a *zero-one matrix*
  - with a *directed graph*




reflexive



irreflexive



symmetric



antisymmetric

# Closures of Relations

- Let  $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$  on  $A = \{1, 2, 3\}$ .



# Closures of Relations

- Let  $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$  on  $A = \{1, 2, 3\}$ .

Is this relation  $R$  reflexive?



# Closures of Relations

- Let  $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$  on  $A = \{1, 2, 3\}$ .

Is this relation  $R$  reflexive?

No. Why?

# Closures of Relations

- Let  $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$  on  $A = \{1, 2, 3\}$ .

Is this relation  $R$  reflexive?

No. Why?  $(2, 2)$  and  $(3, 3)$  are not in  $R$ .





# Closures of Relations

- Let  $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$  on  $A = \{1, 2, 3\}$ .

Is this relation  $R$  reflexive?

No. Why?  $(2, 2)$  and  $(3, 3)$  are not in  $R$ .

The question is what is the minimal relation  $S \supseteq R$  that is reflexive?



# Closures of Relations

- Let  $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$  on  $A = \{1, 2, 3\}$ .

Is this relation  $R$  reflexive?

No. Why?  $(2, 2)$  and  $(3, 3)$  are not in  $R$ .

The question is what is the minimal relation  $S \supseteq R$  that is reflexive?

How to make  $R$  reflexive by minimum number of additions?



# Closures of Relations

- Let  $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$  on  $A = \{1, 2, 3\}$ .

Is this relation  $R$  reflexive?

No. Why?  $(2, 2)$  and  $(3, 3)$  are not in  $R$ .

The question is what is the minimal relation  $S \supseteq R$  that is reflexive?

How to make  $R$  reflexive by minimum number of additions?

Add  $(2, 2)$  and  $(3, 3)$

Then  $S = \{(1, 1), (1, 2), (2, 1), (3, 2), (2, 2), (3, 3)\} \supseteq R$



# Closures of Relations

- Let  $R = \{(1, 1), (1, 2), (2, 1), (3, 2)\}$  on  $A = \{1, 2, 3\}$ .

Is this relation  $R$  reflexive?

No. Why?  $(2, 2)$  and  $(3, 3)$  are not in  $R$ .

The question is what is the minimal relation  $S \supseteq R$  that is reflexive?

How to make  $R$  reflexive by minimum number of additions?

Add  $(2, 2)$  and  $(3, 3)$

Then  $S = \{(1, 1), (1, 2), (2, 1), (3, 2), (2, 2), (3, 3)\} \supseteq R$

The minimal set  $S \supseteq R$  is called *the reflexive closure of  $R$* .



# Reflexive Closure

- The set  $S$  is called *the reflexive closure of  $R$*  if it:



# Reflexive Closure

- The set  $S$  is called *the reflexive closure of  $R$*  if it:
  - ◇ contains  $R$
  - ◇ is reflexive
  - ◇ is minimal (is contained in every reflexive relation  $Q$  that contains  $R$  ( $R \subseteq Q$ ), i.e.,  $S \subseteq Q$ )



# Closures on Relations

- Relations can have different properties:
  - reflexive
  - symmetric
  - transitive



# Closures on Relations

- Relations can have different properties:
  - reflexive
  - symmetric
  - transitive

We define:

- reflexive closures
- symmetric closures
- transitive closures



# Closures

- **Definition** Let  $R$  be a relation on a set  $A$ . A relation  $S$  on  $A$  with property  $P$  is called *the closure of  $R$  with respect to  $P$*  if  $S$  is subset of every relation  $Q$  ( $S \subseteq Q$ ) with property  $P$  that contains  $R$  ( $R \subseteq Q$ ).



# Closures

- **Definition** Let  $R$  be a relation on a set  $A$ . A relation  $S$  on  $A$  with property  $P$  is called *the closure of  $R$  with respect to  $P$*  if  $S$  is subset of every relation  $Q$  ( $S \subseteq Q$ ) with property  $P$  that contains  $R$  ( $R \subseteq Q$ ).

$S$  is the **minimal set** containing  $R$  satisfying the property  $P$ .

# Closures

- **Definition** Let  $R$  be a relation on a set  $A$ . A relation  $S$  on  $A$  with property  $P$  is called *the closure of  $R$  with respect to  $P$*  if  $S$  is subset of every relation  $Q$  ( $S \subseteq Q$ ) with property  $P$  that contains  $R$  ( $R \subseteq Q$ ).

$S$  is the **minimal set** containing  $R$  satisfying the property  $P$ .

**Example:** (symmetric closure)

$$R = \{(1, 2), (1, 3), (2, 2)\} \text{ on } A = \{1, 2, 3\}$$

What is the symmetric closure  $S$  of  $R$ ?

# Closures

- **Definition** Let  $R$  be a relation on a set  $A$ . A relation  $S$  on  $A$  with property  $P$  is called *the closure of  $R$  with respect to  $P$*  if  $S$  is subset of every relation  $Q$  ( $S \subseteq Q$ ) with property  $P$  that contains  $R$  ( $R \subseteq Q$ ).

$S$  is the **minimal set** containing  $R$  satisfying the property  $P$ .

**Example:** (symmetric closure)

$$R = \{(1, 2), (1, 3), (2, 2)\} \text{ on } A = \{1, 2, 3\}$$

What is the symmetric closure  $S$  of  $R$ ?

$$R = \{(1, 2), (1, 3), (2, 2)\} \cup \{(2, 1), (3, 1)\}$$



# Closures

- **Definition** Let  $R$  be a relation on a set  $A$ . A relation  $S$  on  $A$  with property  $P$  is called *the closure of  $R$  with respect to  $P$*  if  $S$  is subset of every relation  $Q$  ( $S \subseteq Q$ ) with property  $P$  that contains  $R$  ( $R \subseteq Q$ ).

$S$  is the **minimal set** containing  $R$  satisfying the property  $P$ .

**Example:** (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is  $R$  transitive?



# Closures

- **Definition** Let  $R$  be a relation on a set  $A$ . A relation  $S$  on  $A$  with property  $P$  is called *the closure of  $R$  with respect to  $P$*  if  $S$  is subset of every relation  $Q$  ( $S \subseteq Q$ ) with property  $P$  that contains  $R$  ( $R \subseteq Q$ ).

$S$  is the **minimal set** containing  $R$  satisfying the property  $P$ .

**Example:** (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is  $R$  transitive? **No**

How to make it transitive?



# Closures

- **Definition** Let  $R$  be a relation on a set  $A$ . A relation  $S$  on  $A$  with property  $P$  is called *the closure of  $R$  with respect to  $P$*  if  $S$  is subset of every relation  $Q$  ( $S \subseteq Q$ ) with property  $P$  that contains  $R$  ( $R \subseteq Q$ ).

$S$  is the **minimal set** containing  $R$  satisfying the property  $P$ .

**Example:** (transitive closure)

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Is  $R$  transitive?      **No**

How to make it transitive?

$$S = \{(1, 2), (2, 2), (2, 3)\} \cup \{(1, 3)\}$$



# Transitive Closure

- We can represent the relation on the graph.





# Transitive Closure

- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path



# Transitive Closure

- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path

**Example:**

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure:  $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



# Transitive Closure


- We can represent the relation on the graph.

Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path

**Example:**

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

Transitive closure:  $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



# Transitive Closure


- We can represent the relation on the graph.

Finding a **transitive closure** corresponds to **finding all pairs** of elements that are connected **with a directed path**

**Example:**

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

**Transitive closure:**  $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



# Transitive Closure



- We can represent the relation on the graph.

Finding a **transitive closure** corresponds to **finding all pairs** of elements that are connected **with a directed path**

**Example:**

$$R = \{(1, 2), (2, 2), (2, 3)\} \text{ on } A = \{1, 2, 3\}$$

**Transitive closure:**  $S = \{(1, 2), (2, 2), (2, 3), (1, 3)\}$



# Paths in Directed Graphs

- **Definition** A *path* from  $a$  to  $b$  in the directed graph  $G$  is a *sequence of edges*  $(x_0, x_1), (x_1, x_2), \dots, (x_{n-1}, x_n)$  in  $G$ , where  $n$  is nonnegative and  $x_0 = a$  and  $x_n = b$ . A path of length  $n \geq 1$  that begins and ends at the same vertex is called a *circuit* or *cycle*.



# Paths in Directed Graphs

- **Definition** A *path* from  $a$  to  $b$  in the directed graph  $G$  is a *sequence of edges*  $(x_0, x_1), (x_1, x_2), \dots, (x_{n-1}, x_n)$  in  $G$ , where  $n$  is nonnegative and  $x_0 = a$  and  $x_n = b$ . A path of length  $n \geq 1$  that begins and ends at the same vertex is called a *circuit* or *cycle*.

**Theorem** Let  $R$  be relation on a set  $A$ . There is a path of length  $n$  from  $a$  to  $b$  if and only if  $(a, b) \in R^n$ .



# Path Length

- **Theorem** Let  $R$  be relation on a set  $A$ . There is a path of length  $n$  from  $a$  to  $b$  if and only if  $(a, b) \in R^n$ .

**Proof** (by induction)





# Path Length

- **Theorem** Let  $R$  be relation on a set  $A$ . There is a path of length  $n$  from  $a$  to  $b$  if and only if  $(a, b) \in R^n$ .

**Proof** (by induction)



**Path of length 1**


# Path Length

- **Theorem** Let  $R$  be relation on a set  $A$ . There is a path of length  $n$  from  $a$  to  $b$  if and only if  $(a, b) \in R^n$ .

**Proof** (by induction)



**Path of length 1**



**Path of length n+1**

# Connectivity Relation

- **Definition** Let  $R$  be a relation on a set  $A$ . The *connectivity relation*  $R^*$  consists of **all pairs**  $(a, b)$  s.t. there is a path (**of any length**) between  $a$  and  $b$  in  $R$ .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$



# Connectivity Relation


- **Definition** Let  $R$  be a relation on a set  $A$ . The *connectivity relation*  $R^*$  consists of **all pairs**  $(a, b)$  s.t. there is a path (**of any length**) between  $a$  and  $b$  in  $R$ .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

**Example:**

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$



# Connectivity Relation

- **Definition** Let  $R$  be a relation on a set  $A$ . The *connectivity relation*  $R^*$  consists of **all pairs**  $(a, b)$  s.t. there is a path (**of any length**) between  $a$  and  $b$  in  $R$ .


$$R^* = \bigcup_{k=1}^{\infty} R^k$$

**Example:**

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$



# Connectivity Relation

- **Definition** Let  $R$  be a relation on a set  $A$ . The *connectivity relation*  $R^*$  consists of **all pairs**  $(a, b)$  s.t. there is a path (**of any length**) between  $a$  and  $b$  in  $R$ .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$


**Example:**

$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$



# Connectivity Relation

- **Definition** Let  $R$  be a relation on a set  $A$ . The *connectivity relation*  $R^*$  consists of **all pairs**  $(a, b)$  s.t. there is a path (**of any length**) between  $a$  and  $b$  in  $R$ .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

**Example:**


$$A = \{1, 2, 3, 4\}$$

$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$

$$R^4 = \emptyset$$



# Connectivity Relation

- **Definition** Let  $R$  be a relation on a set  $A$ . The *connectivity relation*  $R^*$  consists of **all pairs**  $(a, b)$  s.t. there is a path (**of any length**) between  $a$  and  $b$  in  $R$ .

$$R^* = \bigcup_{k=1}^{\infty} R^k$$

**Example:**

$$A = \{1, 2, 3, 4\}$$


$$R = \{(1, 2), (1, 4), (2, 3), (3, 4)\}$$

$$R^2 = \{(1, 3), (2, 4)\}$$

$$R^3 = \{(1, 4)\}$$

$$R^4 = \emptyset$$

$$R^* = \{(1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4)\}$$





# Connectivity

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .



# Connectivity

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

**Proof** (by intuition)

# Connectivity

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

**Proof** (by intuition)

- ◇ There are at most  $n$  different elements we can visit on a path if the path does not have loops



# Connectivity


- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

**Proof** (by intuition)

- ◇ There are at most  $n$  different elements we can visit on a path if the path does not have loops



- ◇ Loops may increase the length but the same node is visited more than once



# Connectivity


- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

**Proof** (by intuition)

- ◇ There are at most  $n$  different elements we can visit on a path if the path does not have loops



- ◇ Loops may increase the length but the same node is visited more than once



# Connectivity

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

$$R^* = \bigcup_{k=1}^n R^k$$



# Connectivity

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

$$R^* = \bigcup_{k=1}^n R^k$$

**Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .



# Connectivity

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

$$R^* = \bigcup_{k=1}^n R^k$$

**Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .

**Recall** Finding a transitive closure corresponds to finding all pairs of elements that are connected with a directed path





# Connectivity

- **Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .



# Connectivity

- **Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .

**Proof**



- **Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .

## Proof

1.  $R^*$  is transitive
2.  $R^* \subseteq S$  whenever  $S$  is a transitive relation containing  $R$

- **Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .

## Proof

1.  $R^*$  is transitive
2.  $R^* \subseteq S$  whenever  $S$  is a transitive relation containing  $R$

1. If  $(a, b) \in R^*$  and  $(b, c) \in R^*$ , then there are paths from  $a$  to  $b$  and from  $b$  to  $c$  in  $R$ . Thus, there is a path from  $a$  to  $c$  in  $R$ . This means that  $(a, c) \in R^*$ .



- **Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .

## Proof

1.  $R^*$  is transitive
  2.  $R^* \subseteq S$  whenever  $S$  is a transitive relation containing  $R$
2. Suppose that  $S$  is a transitive relation containing  $R$ .



# Connectivity

- **Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .

## Proof

1.  $R^*$  is transitive
  2.  $R^* \subseteq S$  whenever  $S$  is a transitive relation containing  $R$
2. Suppose that  $S$  is a transitive relation containing  $R$ .  
Then  $S^n$  is also transitive and  $S^n \subseteq S$ . Why?

- **Theorem:** The transitive closure of a relation  $R$  equals the connectivity relation  $R^*$ .

## Proof

1.  $R^*$  is transitive
2.  $R^* \subseteq S$  whenever  $S$  is a transitive relation containing  $R$

2. Suppose that  $S$  is a transitive relation containing  $R$ .

Then  $S^n$  is also transitive and  $S^n \subseteq S$ . Why?

We have  $S^* \subseteq S$ . Thus,  $R^* \subseteq S^* \subseteq S$



# Find Transitive Closure

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .





# Find Transitive Closure

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

$$\mathbf{M}_{R^*} = \mathbf{M}_R \vee \mathbf{M}_R^{[2]} \vee \mathbf{M}_R^{[3]} \vee \dots \vee \mathbf{M}_R^{[n]}$$

# Find Transitive Closure

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

$$\mathbf{M}_{R^*} = \mathbf{M}_R \vee \mathbf{M}_R^{[2]} \vee \mathbf{M}_R^{[3]} \vee \dots \vee \mathbf{M}_R^{[n]}$$

**Example**

$$\mathbf{M}_R = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$$

$$\mathbf{M}_{R^*} = ?$$



# Simple Transitive Closure Algorithm

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

**procedure** transClosure ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)

// computes  $R^*$  with zero-one matrices

$A := B := \mathbf{M}_R$ ;

**for**  $i := 2$  to  $n$

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

**return**  $B$

//  $B$  is the zero-one matrix for  $R^*$

# Simple Transitive Closure Algorithm

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

**procedure** transClosure ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)

// computes  $R^*$  with zero-one matrices

$A := B := \mathbf{M}_R$ ;

**for**  $i := 2$  to  $n$

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

**return**  $B$

//  $B$  is the zero-one matrix for  $R^*$

This algorithm takes  $\Theta(n^4)$  time.

# Simple Transitive Closure Algorithm

- **Lemma:** Let  $A$  be a set with  $n$  elements, and  $R$  a relation on  $A$ . If there is a path from  $a$  to  $b$  with  $a \neq b$ , then there exists a path of length  $\leq n - 1$ .

**procedure** transClosure ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)

// computes  $R^*$  with zero-one matrices

$A := B := \mathbf{M}_R$ ;

**for**  $i := 2$  to  $n$

$A := A \odot \mathbf{M}_R$

$B := B \vee A$

**return**  $B$

//  $B$  is the zero-one matrix for  $R^*$

This algorithm takes  $\Theta(n^4)$  time.      Why?

# Roy-Warshall Algorithm

```
procedure Warshall ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)
// computes  $R^*$  with zero-one matrices
 $W := \mathbf{M}_R$ ;
for  $k := 1$  to  $n$ 
  for  $i := 1$  to  $n$ 
    for  $j := 1$  to  $n$ 
       $w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$ 
return  $W$ 
//  $W$  is the zero-one matrix for  $R^*$ 
```

# Roy-Warshall Algorithm

**procedure** Warshall ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)

// computes  $R^*$  with zero-one matrices

$W := \mathbf{M}_R$ ;

**for**  $k := 1$  to  $n$

**for**  $i := 1$  to  $n$

**for**  $j := 1$  to  $n$

$w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$

**return**  $W$

//  $W$  is the zero-one matrix for  $R^*$

$w_{ij} = 1$  means there is a path from  $i$  to  $j$  going only through nodes  $\leq k$ .

$$W_{ij}^{[k]} = W_{ij}^{[k-1]} \vee \left( W_{ik}^{[k-1]} \wedge W_{kj}^{[k-1]} \right)$$

# Roy-Warshall Algorithm

```
procedure Warshall ( $\mathbf{M}_R$ : zero-one  $n \times n$  matrix)  
// computes  $R^*$  with zero-one matrices  
 $W := \mathbf{M}_R$ ;  
for  $k := 1$  to  $n$   
  for  $i := 1$  to  $n$   
    for  $j := 1$  to  $n$   
       $w_{ij} := w_{ij} \vee (w_{ik} \wedge w_{kj})$   
return  $W$   
//  $W$  is the zero-one matrix for  $R^*$ 
```

$w_{ij} = 1$  means there is a path from  $i$  to  $j$  going only through nodes  $\leq k$ .

$$W_{ij}^{[k]} = W_{ij}^{[k-1]} \vee \left( W_{ik}^{[k-1]} \wedge W_{kj}^{[k-1]} \right)$$


This algorithm takes  $\Theta(n^3)$  time.





# Example


Find the matrices  $W_0$ ,  $W_1$ ,  $W_2$ ,  $W_3$ , and  $W_4$ . The matrix  $W_4$  is the **transitive closure** of  $R$ .



Let  $v_1 = a$ ,  $v_2 = b$ ,  $v_3 = c$ ,  $v_4 = d$ .

# Example

Find the matrices  $W_0, W_1, W_2, W_3$ , and  $W_4$ . The matrix  $W_4$  is the **transitive closure** of  $R$ .




Let  $v_1 = a, v_2 = b, v_3 = c, v_4 = d$ .

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

# Example

Find the matrices  $W_0, W_1, W_2, W_3$ , and  $W_4$ . The matrix  $W_4$  is the **transitive closure** of  $R$ .




Let  $v_1 = a, v_2 = b, v_3 = c, v_4 = d$ .

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

# Example

Find the matrices  $W_0, W_1, W_2, W_3$ , and  $W_4$ . The matrix  $W_4$  is the **transitive closure** of  $R$ .



Let  $v_1 = a, v_2 = b, v_3 = c, v_4 = d$ .


$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \mathbf{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \mathbf{1} \\ 1 & 0 & 0 & 1 \\ \mathbf{1} & 0 & 1 & \mathbf{1} \end{bmatrix}$$

# Example

Find the matrices  $W_0, W_1, W_2, W_3$ , and  $W_4$ . The matrix  $W_4$  is the **transitive closure** of  $R$ .



Let  $v_1 = a, v_2 = b, v_3 = c, v_4 = d$ .

$$W_0 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_2 = W_1 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

$$W_3 = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & 0 & 1 \\ \textcolor{red}{1} & 0 & 1 & \textcolor{red}{1} \end{bmatrix}$$

$$W_4 = \begin{bmatrix} \textcolor{red}{1} & 0 & \textcolor{red}{1} & 1 \\ 1 & 0 & 1 & \textcolor{red}{1} \\ 1 & 0 & \textcolor{red}{1} & 1 \\ \textcolor{red}{1} & 0 & 1 & \textcolor{red}{1} \end{bmatrix}$$

# $n$ -ary Relations

- **Definition** An  $n$ -ary relation  $R$  on sets  $A_1, \dots, A_n$ , written as  $R : A_1, \dots, A_n$ , is a subset  $R \subseteq A_1 \times \dots \times A_n$ .



# $n$ -ary Relations

- **Definition** An  $n$ -ary relation  $R$  on sets  $A_1, \dots, A_n$ , written as  $R : A_1, \dots, A_n$ , is a subset  $R \subseteq A_1 \times \dots \times A_n$ .
  - The sets  $A_i$ 's are called the *domains* of  $R$ .

# $n$ -ary Relations

- **Definition** An  $n$ -ary relation  $R$  on sets  $A_1, \dots, A_n$ , written as  $R : A_1, \dots, A_n$ , is a subset  $R \subseteq A_1 \times \dots \times A_n$ .
  - The sets  $A_i$ 's are called the *domains* of  $R$ .
  - The *degree* of  $R$  is  $n$ .





# $n$ -ary Relations

- **Definition** An  $n$ -ary relation  $R$  on sets  $A_1, \dots, A_n$ , written as  $R : A_1, \dots, A_n$ , is a subset  $R \subseteq A_1 \times \dots \times A_n$ .
  - The sets  $A_i$ 's are called the *domains* of  $R$ .
  - The *degree* of  $R$  is  $n$ .
  - $R$  is *functional* in domain  $A_i$  if it contains **at most one**  $n$ -tuple  $(\dots, a_i, \dots)$  for any value  $a_i$  within domain  $A_i$ .



# Relational Databases

- A *relational database* is essentially an  $n$ -ary relation  $R$ .



# Relational Databases

- A *relational database* is essentially an *n*-ary relation  $R$ .
- A domain  $A_i$  is a *primary key* for the database if the relation  $R$  is *functional* in  $A_i$ .




# Relational Databases

- A *relational database* is essentially an *n*-ary relation *R*.
- A domain  $A_i$  is a *primary key* for the database if the relation *R* is *functional* in  $A_i$ .
- A *composite key* for the database is a set of domains  $\{A_i, A_j, \dots\}$  such that *R* contains **at most 1 *n*-tuple**  $(\dots, a_i, \dots, a_j, \dots)$  for each composite value  $(a_i, a_j, \dots) \in A_i \times A_j \times \dots$ .

# Relational Databases

## *E-R Diagram*



# Selection Operators

- Let  $A$  be any  $n$ -ary domain  $A = A_1 \times \cdots \times A_n$ , and let  $C : A \rightarrow \{T, F\}$  be any *condition* (predicate) on elements ( $n$ -tuples) of  $A$ .



# Selection Operators

- Let  $A$  be any  *$n$ -ary domain*  $A = A_1 \times \cdots \times A_n$ , and let  $C : A \rightarrow \{T, F\}$  be any *condition* (predicate) on elements ( $n$ -tuples) of  $A$ .
- The *selection operator*  $s_C$  is the operator that maps any ( $n$ -ary) relation  $R$  on  $A$  to the  $n$ -ary relation of all  $n$ -tuples from  $R$  that *satisfy*  $C$ .

# Selection Operators

- Let  $A$  be any  *$n$ -ary domain*  $A = A_1 \times \cdots \times A_n$ , and let  $C : A \rightarrow \{T, F\}$  be any *condition* (predicate) on elements ( $n$ -tuples) of  $A$ .
- The *selection operator*  $s_C$  is the operator that maps any ( $n$ -ary) relation  $R$  on  $A$  to the  $n$ -ary relation of all  $n$ -tuples from  $R$  that *satisfy*  $C$ .

$$- \forall R \subseteq A,$$

$$\begin{aligned} s_C(R) &= R \cap \{a \in A \mid s_C(a) = T\} \\ &= \{a \in R \mid s_C(a) = T\}. \end{aligned}$$





# Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$



# Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$

- Suppose that we have a domain

$$\begin{aligned} &\textit{UpperLevel}(\textit{name}, \textit{standing}, \textit{ssn}) \\ &::= [(\textit{standing} = \textit{junior}) \vee (\textit{standing} = \textit{senior})] \end{aligned}$$



# Selection Operator Example

- Suppose that we have a domain

$$A = \textit{StudentName} \times \textit{Standing} \times \textit{SocSecNos}$$

- Suppose that we have a domain

$$\begin{aligned} &\textit{UpperLevel}(\textit{name}, \textit{standing}, \textit{ssn}) \\ &:\equiv [(\textit{standing} = \textit{junior}) \vee (\textit{standing} = \textit{senior})] \end{aligned}$$

- Then,  $\textit{SUpperLevel}$  is the selection operator that takes any relation  $R$  on  $A$  (database of students) and produces a relation consisting of just the upper-level classes (juniors and seniors).



# Projection Operators

- Let  $A = A_1 \times \cdots \times A_n$  be any  $n$ -ary domain, and let  $\{i_k\} = (i_1, \dots, i_m)$  be a sequence of indices all falling in the range 1 to  $n$ .  
i.e., where  $1 \leq i_k \leq n$  for all  $1 \leq k \leq m$ .



# Projection Operators

- Let  $A = A_1 \times \cdots \times A_n$  be any  *$n$ -ary domain*, and let  $\{i_k\} = (i_1, \dots, i_m)$  be a sequence of indices all falling in the range 1 to  $n$ .

i.e., where  $1 \leq i_k \leq n$  for all  $1 \leq k \leq m$ .

- Then the *projection operator* on  $n$ -tuples

$$P_{\{i_k\}} : A \rightarrow A_{i_1} \times \cdots \times A_{i_m}$$

is defined by

$$P_{\{i_k\}}(a_1, \dots, a_n) = (a_{i_1}, \dots, a_{i_m})$$



# Projection Example

- Suppose that we have a tenary domain

$$Cars = Model \times Year \times Color \ (n = 3)$$



# Projection Example

- Suppose that we have a tenary domain

$$\textit{Cars} = \textit{Model} \times \textit{Year} \times \textit{Color} \quad (n = 3)$$

- Consider the index sequence  $\{i_k\} = \{1, 3\}$  ( $m = 2$ )



# Projection Example

- Suppose that we have a ternary domain

$$Cars = Model \times Year \times Color \quad (n = 3)$$

- Consider the index sequence  $\{i_k\} = \{1, 3\}$  ( $m = 2$ )

- Then the projection  $P_{\{i_k\}}$  simply maps each tuple  $(a_1, a_2, a_3) = (model, year, color)$  to its image:

$$(a_{i_1}, a_{i_2}) = (a_1, a_3) = (model, color)$$





# Projection Example

- Suppose that we have a tenary domain

$$Cars = Model \times Year \times Color \quad (n = 3)$$

- Consider the index sequence  $\{i_k\} = \{1, 3\}$  ( $m = 2$ )

- Then the projection  $P_{\{i_k\}}$  simply maps each tuple  $(a_1, a_2, a_3) = (model, year, color)$  to its image:

$$(a_{i_1}, a_{i_2}) = (a_1, a_3) = (model, color)$$

- This operator can be usefully applied to a whole relation  $R \subseteq Cars$  (database of cars) to obtain a list of *model/color* combinations available.



# Join Operator

- Puts two relations together to form a sort of *combined relation*.



# Join Operator

- Puts two relations together to form a sort of *combined relation*.
- If the tuple  $(A, B)$  appears in  $R_1$ , and the tuple  $(B, C)$  appears in  $R_2$ , then the tuple  $(A, B, C)$  appears in the *join*  $J(R_1, R_2)$ .



# Join Operator

- Puts two relations together to form a sort of *combined relation*.
- If the tuple  $(A, B)$  appears in  $R_1$ , and the tuple  $(B, C)$  appears in  $R_2$ , then the tuple  $(A, B, C)$  appears in the *join*  $J(R_1, R_2)$ .
- $A, B, C$  can also be sequences of elements rather than single elements.



# Join Example

- Suppose that  $R_1$  is a teaching assignment table, relating *Professors* to *Courses*.



# Join Example

- Suppose that  $R_1$  is a teaching assignment table, relating *Professors* to *Courses*.
- Suppose that  $R_2$  is a room assignment table relating *Courses* to *Rooms* and *Times*.



# Join Example

- Suppose that  $R_1$  is a teaching assignment table, relating *Professors* to *Courses*.
- Suppose that  $R_2$  is a room assignment table relating *Courses* to *Rooms* and *Times*.
- Then  $J(R_1, R_2)$  is like your **class schedule**, listing *(professor, course, room, time)*.



# Next Lecture

- relation III ...

