



# CS215 DISCRETE MATH

Dr. QI WANG

Department of Computer Science and Engineering

Office: Room413, CoE South Tower

Email: [wangqi@sustech.edu.cn](mailto:wangqi@sustech.edu.cn)

# Application of Number Theory

- G. H. Hardy (1877 - 1947)

In his 1940 autobiography *A Mathematician's Apology*, Hardy wrote “The great modern achievements of applied mathematics have been in *relativity* and *quantum mechanics*, and these subjects are, at present, **almost as ‘useless’ as the theory of numbers.**”



# Application of Number Theory

- G. H. Hardy (1877 - 1947)

In his 1940 autobiography *A Mathematician's Apology*, Hardy wrote “The great modern achievements of applied mathematics have been in relativity and quantum mechanics, and these subjects are, at present, **almost as ‘useless’ as the theory of numbers.**”



If he could see the world now, Hardy would be spinning in his grave.

# Number Theory

- *Number theory* is a branch of mathematics that explores integers and their properties, is the basis of **cryptography**, **coding theory**, **computer security**, **e-commerce**, etc.



# Number Theory

- *Number theory* is a branch of mathematics that explores integers and their properties, is the basis of **cryptography**, **coding theory**, **computer security**, **e-commerce**, etc.
- At one point, the largest employer of mathematicians in the United States, and probably the world, was the **National Security Agency** (NSA). The NSA is the largest spy agency in the US (bigger than CIA, Central Intelligence Agency), and has the responsibility for code design and breaking.



# Division

- If  $a$  and  $b$  are integers with  $a \neq 0$ , we say that  $a$  divides  $b$  if there is an integer  $k$  such that  $b = ak$ , or equivalently  $b/a$  is an integer. In this case, we say that  $a$  is a *factor* or *divisor* of  $b$ , and  $b$  is a *multiple* of  $a$ . (We use the notations  $a|b$ ,  $a \nmid b$ )



# Division

- If  $a$  and  $b$  are integers with  $a \neq 0$ , we say that  $a$  divides  $b$  if there is an integer  $k$  such that  $b = ak$ , or equivalently  $b/a$  is an integer. In this case, we say that  $a$  is a *factor* or *divisor* of  $b$ , and  $b$  is a *multiple* of  $a$ . (We use the notations  $a|b$ ,  $a \nmid b$ )

## Example

◇  $4 \mid 24$

◇  $3 \nmid 7$



# Divisibility

- **All integers divisible by  $d > 0$  can be enumerated as:**  
 $\dots, -kd, \dots, -2d, -d, 0, d, 2d, \dots, kd, \dots$





# Divisibility

- **All integers divisible by  $d > 0$**  can be **enumerated** as:  
$$\dots, -kd, \dots, -2d, -d, 0, d, 2d, \dots, kd, \dots$$
- **Question:** Let  $n$  and  $d$  be two positive integers. How many positive integers **not exceeding  $n$**  are divisible by  $d$ ?



# Divisibility

- **All integers divisible by  $d > 0$**  can be **enumerated** as:  
 $\dots, -kd, \dots, -2d, -d, 0, d, 2d, \dots, kd, \dots$
- **Question:** Let  $n$  and  $d$  be two positive integers. How many positive integers **not exceeding  $n$**  are divisible by  $d$ ?

**Answer:** Count the number of integers such that  $0 < kd \leq n$ . Therefore, there are  $\lfloor n/d \rfloor$  such positive integers.



# Divisibility

## ■ Properties

Let  $a, b, c$  be integers. Then the following hold:

- (i) if  $a|b$  and  $a|c$ , then  $a|(b + c)$
- (ii) if  $a|b$  then  $a|bc$  for all integers  $c$
- iii) if  $a|b$  and  $b|c$ , then  $a|c$



# Divisibility

## ■ Properties

Let  $a, b, c$  be integers. Then the following hold:

- (i) if  $a|b$  and  $a|c$ , then  $a|(b + c)$
- (ii) if  $a|b$  then  $a|bc$  for all integers  $c$
- iii) if  $a|b$  and  $b|c$ , then  $a|c$

**Proof.**



# Divisibility

- **Corollary** If  $a, b, c$  are integers, where  $a \neq 0$ , such that  $a|b$  and  $a|c$ , then  $a|(mb + nc)$  whenever  $m$  and  $n$  are integers.



# Divisibility

- **Corollary** If  $a, b, c$  are integers, where  $a \neq 0$ , such that  $a|b$  and  $a|c$ , then  $a|(mb + nc)$  whenever  $m$  and  $n$  are integers.

**Proof.** By part (ii) and part (i) of Properties.

# The Division Algorithm

- If  $a$  is an integer and  $d$  a positive integer, then there are **unique** integers  $q$  and  $r$ , with  $0 \leq r < d$ , such that  $a = dq + r$ . In this case,  $d$  is called the *divisor*,  $a$  is called the *dividend*,  $q$  is called the *quotient*, and  $r$  is called the *remainder*.



# The Division Algorithm

- If  $a$  is an integer and  $d$  a positive integer, then there are **unique** integers  $q$  and  $r$ , with  $0 \leq r < d$ , such that  $a = dq + r$ . In this case,  $d$  is called the *divisor*,  $a$  is called the *dividend*,  $q$  is called the *quotient*, and  $r$  is called the *remainder*.

In this case, we use the notations  $q = a \text{ div } d$  and  $r = a \bmod d$ .





# Congruence Relation

- If  $a$  and  $b$  are integers and  $m$  is a positive integer, then  $a$  is *congruent to  $b$  modulo  $m$  if  $m$  divides  $a - b$* , denoted by  $a \equiv b \pmod{m}$ . This is called *congruence* and  $m$  is its *modulus*.



# Congruence Relation

- If  $a$  and  $b$  are integers and  $m$  is a positive integer, then  $a$  is *congruent to  $b$  modulo  $m$  if  $m$  divides  $a - b$* , denoted by  $a \equiv b \pmod{m}$ . This is called *congruence* and  $m$  is its *modulus*.

## Example

- ◇  $15 \equiv 3 \pmod{6}$
- ◇  $-1 \equiv 11 \pmod{6}$



# More on Congruences

- Let  $m$  be a positive integer. The integers  $a$  and  $b$  are congruent modulo  $m$  if and only if there is an integer  $k$  such that  $a = b + km$ .



# More on Congruences

- Let  $m$  be a positive integer. The integers  $a$  and  $b$  are congruent modulo  $m$  if and only if there is an integer  $k$  such that  $a = b + km$ .

**Proof.**

“only if” part

“if” part



# $(\bmod m)$ and $\bmod m$ Notations

- $a \equiv b \pmod{m}$  and  $a \bmod m = b$  are different.
  - ◇  $a \equiv b \pmod{m}$  is a **relation** on the set of integers
  - ◇ In  $a \bmod m = b$ , the notation **mod** denotes a function



# $(\bmod m)$ and $\bmod m$ Notations

- $a \equiv b \pmod{m}$  and  $a \bmod m = b$  are different.
  - ◇  $a \equiv b \pmod{m}$  is a **relation** on the set of integers
  - ◇ In  $a \bmod m = b$ , the notation **mod** denotes a function
- Let  $a$  and  $b$  be integers, and let  $m$  be a positive integer.  
Then  $a \equiv b \bmod m$  if and only if  $a \bmod m = b \bmod m$



# $(\text{mod } m)$ and $\text{mod } m$ Notations

- $a \equiv b \pmod{m}$  and  $a \bmod m = b$  are different.
  - ◇  $a \equiv b \pmod{m}$  is a **relation** on the set of integers
  - ◇ In  $a \bmod m = b$ , the notation **mod** denotes a function
- Let  $a$  and  $b$  be integers, and let  $m$  be a positive integer.  
Then  $a \equiv b \bmod m$  if and only if  $a \bmod m = b \bmod m$

**Proof.**



# Congruences of Sums and Products

- Let  $m$  be a positive integer. If  $a \equiv b \pmod{m}$  and  $c \equiv d \pmod{m}$ , then  $a + c \equiv b + d \pmod{m}$  and  $ac \equiv bd \pmod{m}$





# Congruences of Sums and Products

- Let  $m$  be a positive integer. If  $a \equiv b \pmod{m}$  and  $c \equiv d \pmod{m}$ , then  $a + c \equiv b + d \pmod{m}$  and  $ac \equiv bd \pmod{m}$

**Proof.**

# Algebraic Manipulation of Congruences

- If  $a \equiv b \pmod{m}$ , then
  - $c \cdot a \equiv c \cdot b \pmod{m}$ ?
  - $c + a \equiv c + b \pmod{m}$ ?
  - $a/c \equiv b/c \pmod{m}$ ?



# Algebraic Manipulation of Congruences

- If  $a \equiv b \pmod{m}$ , then
  - $c \cdot a \equiv c \cdot b \pmod{m}$ ?
  - $c + a \equiv c + b \pmod{m}$ ?
  - $a/c \equiv b/c \pmod{m}$ ?

$$14 \equiv 8 \pmod{6} \text{ but } 7 \not\equiv 4 \pmod{6}$$



# Computing the mod Function

- **Corollary** Let  $m$  be a positive integer and let  $a$  and  $b$  be integers. Then

$$(a + b) \bmod m = ((a \bmod m) + (b \bmod m)) \bmod m$$

$$ab \bmod m = ((a \bmod m)(b \bmod m)) \bmod m$$



# Computing the mod Function

- **Corollary** Let  $m$  be a positive integer and let  $a$  and  $b$  be integers. Then

$$(a + b) \bmod m = ((a \bmod m) + (b \bmod m)) \bmod m$$
$$ab \bmod m = ((a \bmod m)(b \bmod m)) \bmod m$$

**Proof.**



# Arithmetic Modulo $m$

- Let  $\mathbf{Z}_m$  be the set of nonnegative integers less than  $m$ :  $\{0, 1, \dots, m - 1\}$ .



# Arithmetic Modulo $m$

- Let  $\mathbf{Z}_m$  be the set of nonnegative integers less than  $m$ :  $\{0, 1, \dots, m - 1\}$ .

$$+_m : a +_m b = (a + b) \bmod m$$

$$\cdot_m : a \cdot_m b = ab \bmod m$$



# Arithmetic Modulo $m$

- Let  $\mathbf{Z}_m$  be the set of nonnegative integers less than  $m$ :  $\{0, 1, \dots, m - 1\}$ .

$$+_m : a +_m b = (a + b) \bmod m$$

$$\cdot_m : a \cdot_m b = ab \bmod m$$

## Example

$$\diamond 7 +_{11} 9 = ?$$

$$\diamond 7 \cdot_{11} 9 = ?$$





# Arithmetic Modulo $m$

- **Closure:** if  $a, b \in \mathbf{Z}_m$ , then  $a +_m b, a \cdot_m b \in \mathbf{Z}_m$



# Arithmetic Modulo $m$

- **Closure**: if  $a, b \in \mathbf{Z}_m$ , then  $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity**: if  $a, b, c \in \mathbf{Z}_m$ , then  
 $(a +_m b) +_m c = a +_m (b +_m c)$  and  
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$



# Arithmetic Modulo $m$

- **Closure**: if  $a, b \in \mathbf{Z}_m$ , then  $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity**: if  $a, b, c \in \mathbf{Z}_m$ , then  
 $(a +_m b) +_m c = a +_m (b +_m c)$  and  
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$
- **Identity elements**:  $a +_m 0 = a$  and  $a \cdot_m 1 = a$



# Arithmetic Modulo $m$

- **Closure:** if  $a, b \in \mathbf{Z}_m$ , then  $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity:** if  $a, b, c \in \mathbf{Z}_m$ , then  
 $(a +_m b) +_m c = a +_m (b +_m c)$  and  
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$
- **Identity elements:**  $a +_m 0 = a$  and  $a \cdot_m 1 = a$
- **Additive inverses:** if  $a \neq 0$  and  $a \in \mathbf{Z}_m$ , then  $m - a$  is an additive inverse of  $a$  modulo  $m$



# Arithmetic Modulo $m$

- **Closure**: if  $a, b \in \mathbf{Z}_m$ , then  $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity**: if  $a, b, c \in \mathbf{Z}_m$ , then  
 $(a +_m b) +_m c = a +_m (b +_m c)$  and  
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$
- **Identity elements**:  $a +_m 0 = a$  and  $a \cdot_m 1 = a$
- **Additive inverses**: if  $a \neq 0$  and  $a \in \mathbf{Z}_m$ , then  $m - a$  is an additive inverse of  $a$  modulo  $m$
- **Commutativity**: if  $a, b \in \mathbf{Z}_m$ , then  $a +_m b = b +_m a$



# Arithmetic Modulo $m$

- **Closure**: if  $a, b \in \mathbf{Z}_m$ , then  $a +_m b, a \cdot_m b \in \mathbf{Z}_m$
- **Associativity**: if  $a, b, c \in \mathbf{Z}_m$ , then  
 $(a +_m b) +_m c = a +_m (b +_m c)$  and  
 $(a \cdot_m b) \cdot_m c = a \cdot_m (b \cdot_m c)$
- **Identity elements**:  $a +_m 0 = a$  and  $a \cdot_m 1 = a$
- **Additive inverses**: if  $a \neq 0$  and  $a \in \mathbf{Z}_m$ , then  $m - a$  is an additive inverse of  $a$  modulo  $m$
- **Commutativity**: if  $a, b \in \mathbf{Z}_m$ , then  $a +_m b = b +_m a$
- **Distributivity**: if  $a, b, c \in \mathbf{Z}_m$ , then  
 $a \cdot_m (b +_m c) = (a \cdot_m b) +_m (a \cdot_m c)$  and  
 $(a +_m b) \cdot_m c = (a \cdot_m c) +_m (b \cdot_m c)$



# Group

- A set  $G$  of elements, along with a **binary operation**  $\star$ , must satisfy the following four properties to be called a *group*.



# Group

- A set  $G$  of elements, along with a **binary operation**  $\star$ , must satisfy the following four properties to be called a **group**.

**Closure:** If  $a, b \in G$ , then  $a \star b = c$  also belongs to  $G$ .

**Associativity:**  $(a \star b) \star c = a \star (b \star c)$

**Identity element:** There is a **unique** element  $1_e$ , such that for every  $a \in G$ , we have  $a \star 1_e = a$ .

**Inverse:** For every  $a \in G$ , there exists an element, denoted by  $a^{-1}$ , such that  $a \star a^{-1} = 1_e$ .





# Group

- A set  $G$  of elements, along with a **binary operation**  $\star$ , must satisfy the following four properties to be called a **group**.

**Closure:** If  $a, b \in G$ , then  $a \star b = c$  also belongs to  $G$ .

**Associativity:**  $(a \star b) \star c = a \star (b \star c)$

**Identity element:** There is a **unique** element  $1_e$ , such that for every  $a \in G$ , we have  $a \star 1_e = a$ .

**Inverse:** For every  $a \in G$ , there exists an element, denoted by  $a^{-1}$ , such that  $a \star a^{-1} = 1_e$ .

**Example:**

$(\mathbb{Z}, +)$ ,  $(\mathbb{Q}, +)$ ,  $(\mathbb{R}, +)$ ,  $(\mathbb{M}_{n \times n}, +)$  ?

$(\mathbb{Z}^*, \times)$ ,  $(\mathbb{Q}^*, \times)$ ,  $(\mathbb{R}^*, \times)$ ,  $(\mathbb{M}_{n \times n}^*, \cdot)$  ?



# Permutation Group

- Let  $s_n = \langle 1, 2, \dots, n \rangle$  denote a *sequence* of integers 1 through  $n$ . Denote by  $P_n$  the set of all *permutations* of the sequence  $s_n$ .



# Permutation Group

- Let  $s_n = \langle 1, 2, \dots, n \rangle$  denote a *sequence* of integers 1 through  $n$ . Denote by  $P_n$  the set of all *permutations* of the sequence  $s_n$ .

For example,  $s_3 = \langle 1, 2, 3 \rangle$

$$P_3 = \{ \langle 1, 2, 3 \rangle, \langle 1, 3, 2 \rangle, \langle 2, 1, 3 \rangle, \langle 2, 3, 1 \rangle, \langle 3, 1, 2 \rangle, \langle 3, 2, 1 \rangle \}$$



# Permutation Group

- Let  $s_n = \langle 1, 2, \dots, n \rangle$  denote a *sequence* of integers 1 through  $n$ . Denote by  $P_n$  the set of all *permutations* of the sequence  $s_n$ .

For example,  $s_3 = \langle 1, 2, 3 \rangle$

$$P_3 = \{ \langle 1, 2, 3 \rangle, \langle 1, 3, 2 \rangle, \langle 2, 1, 3 \rangle, \langle 2, 3, 1 \rangle, \langle 3, 1, 2 \rangle, \langle 3, 2, 1 \rangle \}$$

- Define a binary operation  $\circ$  on the elements of  $P_n$ : for  $\rho, \pi \in P_n$ ,  $\pi \circ \rho$  denotes a *re-permutation* of the elements of  $\rho$  according to the elements of  $\pi$ .



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?  
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$



# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?  
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$
- We can verify the other three properties.

$$\rho_1 \circ (\rho_2 \circ \rho_3) = (\rho_1 \circ \rho_2) \circ \rho_3$$

$$\langle 1, 2, 3 \rangle \circ \rho = \rho \circ \langle 1, 2, 3 \rangle = \rho$$

For each  $\rho \in P_3$ , there exists another unique  $\pi \in P_3$  such that

$$\rho \circ \pi = \pi \circ \rho = \langle 1, 2, 3 \rangle$$





# Permutation Group

- Consider  $s_3 = \langle 1, 2, 3 \rangle$ , and  
 $P_3 = \{ \langle p_1, p_2, p_3 \rangle \mid p_1, p_2, p_3 \in s_3 \text{ with } p_1 \neq p_2 \neq p_3 \}$ .
- $\pi = \langle 3, 2, 1 \rangle$ ,  $\rho = \langle 1, 3, 2 \rangle$ , what is  $\pi \circ \rho$ ?  
 $\pi \circ \rho = \langle 2, 3, 1 \rangle \in P_3$
- We can verify the other three properties.  
$$\rho_1 \circ (\rho_2 \circ \rho_3) = (\rho_1 \circ \rho_2) \circ \rho_3$$
$$\langle 1, 2, 3 \rangle \circ \rho = \rho \circ \langle 1, 2, 3 \rangle = \rho$$

For each  $\rho \in P_3$ , there exists another unique  $\pi \in P_3$  such that

$$\rho \circ \pi = \pi \circ \rho = \langle 1, 2, 3 \rangle$$

$(P_n, \circ)$  is called a *permutation group*.



# Abelian Group

- If the operation on the set elements is *commutative*, the group is called an *abelian group*. ( $a \star b = b \star a$ )



# Abelian Group

- If the operation on the set elements is *commutative*, the group is called an *abelian group*. ( $a \star b = b \star a$ )

- **Example**

$(\mathbb{Z}, +)$ ,  $(\mathbb{Q}, +)$ ,  $(\mathbb{R}, +)$ ,  $(\mathbb{M}_{n \times n}, +)$  ?

$(GL(n), \cdot)$ ,  $(P_n, \circ)$  ?



# Abelian Group

- If the operation on the set elements is *commutative*, the group is called an *abelian group*. ( $a \star b = b \star a$ )

- **Example**

$$(\mathbb{Z}, +), (\mathbb{Q}, +), (\mathbb{R}, +), (\mathbb{M}_{n \times n}, +) ?$$

$$(GL(n), \cdot), (P_n, \circ) ?$$

- If the group operation is referred to as *addition* (*multiplication*), then the group also allows for *subtraction* (*division*).

$$a - b = a + (-b)$$

$$a/b = a \cdot b^{-1}$$



# Ring

- If  $(R, +)$  is an *abelian group*, we define one more operation (denoted as *multiplication*  $\times$  for convenience) to have a *ring*  $(R, +, \times)$  satisfying the following properties.



# Ring

- If  $(R, +)$  is an *abelian group*, we define one more operation (denoted as *multiplication*  $\times$  for convenience) to have a *ring*  $(R, +, \times)$  satisfying the following properties.

**Closure:**  $R$  must be closed w.r.t.  $\times$

**Associativity:**  $(a \times b) \times c = a \times (b \times c)$

**Distributivity:**  $a \times (b + c) = a \times b + a \times c$   
 $(a + b) \times c = a \times c + b \times c$



# Ring

- If  $(R, +)$  is an *abelian group*, we define one more operation (denoted as *multiplication*  $\times$  for convenience) to have a *ring*  $(R, +, \times)$  satisfying the following properties.

**Closure:**  $R$  must be closed w.r.t.  $\times$

**Associativity:**  $(a \times b) \times c = a \times (b \times c)$

**Distributivity:**  $a \times (b + c) = a \times b + a \times c$   
 $(a + b) \times c = a \times c + b \times c$

**Example:**

$(\mathbb{Z}, +, \times)$ ,  $(\mathbb{Q}, +, \times)$ ,  $(\mathbb{R}, +, \times)$ ,  $(\mathbb{M}_{n \times n}, +, \cdot)$  ?



# Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ( $ab = ba$ )





# Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ( $ab = ba$ )
- An *integral domain*  $(R, +, \times)$  is a *commutative ring* that satisfies the following two additional properties.

**Identity element** for multiplication:  $a1 = 1a = a$

**Nonzero product** for any two nonzero elements:  
if  $ab = 0$ , then either  $a$  or  $b$  **must be** 0.



# Commutative Ring, Integral Domain

- A *ring* is *commutative* if the **multiplication operation** is *commutative* for all elements in the ring. ( $ab = ba$ )
- An *integral domain*  $(R, +, \times)$  is a *commutative ring* that satisfies the following two additional properties.

**Identity element** for multiplication:  $a1 = 1a = a$

**Nonzero product** for any two nonzero elements:

if  $ab = 0$ , then either  $a$  or  $b$  **must be** 0.

**Example:**

$(\mathbb{Z}, +, \times)$ ,  $(\mathbb{Q}, +, \times)$ ,  $(\mathbb{R}, +, \times)$  ?

$(\mathbb{Z}_m, +, \times)$ ,  $(M_{n \times n}, +, \cdot)$  ?



# Field

- A *field*, denoted by  $(F, +, \times)$ , is an *integral domain* whose elements satisfy the following additional property.

**Inverse** for multiplication: For every  $a \in F$ , there exists an element  $b$ , denoted by  $a^{-1}$ , such that  $ab = ba = 1$ .



# Field

- A *field*, denoted by  $(F, +, \times)$ , is an *integral domain* whose elements satisfy the following additional property.

**Inverse** for multiplication: For every  $a \in F$ , there exists an element  $b$ , denoted by  $a^{-1}$ , such that  $ab = ba = 1$ .

- **Example:**

$(\mathbb{Z}, +, \times)$ ,  $(\mathbb{Q}, +, \times)$ ,  $(\mathbb{R}, +, \times)$  ?

$(\mathbb{Z}_p, +, \times)$  ?



# Representations of Integers

- We may use *decimal* (*base 10*) or *binary* or *octal* or *hexadecimal* or other notations to represent integers.



# Representations of Integers

- We may use *decimal* (*base 10*) or *binary* or *octal* or *hexadecimal* or other notations to represent integers.
- Let  $b > 1$  be an integer. Then if  $n$  is a positive integer, it can be expressed **uniquely in the form**  
$$n = a_k b^k + a_{k-1} b^{k-1} + \dots + a_1 b + a_0,$$
 where  $k$  is nonnegative,  $a_i$ 's are nonnegative integers less than  $b$ . The representation of  $n$  is called *the base- $b$  expansion of  $n$*  and is denoted by  $(a_k a_{k-1} \dots a_1 a_0)_b$ .



# Base- $b$ Expansions

- To get the decimal expansion is easy.



# Base- $b$ Expansions

- To get the decimal expansion is easy.

## Example

- ◇  $(10101111)_2 = 2^8 + 2^6 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 = 351$
- ◇  $(7016)_8 = 7 \cdot 8^3 + 1 \cdot 8 + 6 = 3598$



# Base- $b$ Expansions

- To get the decimal expansion is easy.

## Example

- ◇  $(10101111)_2 = 2^8 + 2^6 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 = 351$
- ◇  $(7016)_8 = 7 \cdot 8^3 + 1 \cdot 8 + 6 = 3598$

- Conversions between binary, octal, hexadecimal expansions are easy.



# Base- $b$ Expansions

- To get the decimal expansion is easy.

## Example

- ◇  $(101011111)_2 = 2^8 + 2^6 + 2^4 + 2^3 + 2^2 + 2^1 + 2^0 = 351$
- ◇  $(7016)_8 = 7 \cdot 8^3 + 1 \cdot 8 + 6 = 3598$

- Conversions between binary, octal, hexadecimal expansions are easy.

## Example

- ◇  $(101011111)_2 = (\underline{101}\overline{011}\underline{111}) = (537)_8$
- ◇  $(7016)_8 = (\underline{111}\overline{000}\underline{001}\overline{110})_2$   
 $= (\underline{111}\overline{000}\underline{001}\overline{110})_2 = (E0E)_{16}$

# Base- $b$ Expansions

$$\begin{aligned}n &= a_k b^k + a_{k-1} b^{k-1} + a_{k-2} b^{k-2} + \cdots + a_2 b^2 + a_1 b + a_0 \\&= b(a_k b^{k-1} + a_{k-1} b^{k-2} + a_{k-2} b^{k-3} + \cdots + a_2 b + a_1) + \textcolor{red}{a_0} \\&= b(b(a_k b^{k-2} + a_{k-1} b^{k-3} + a_{k-2} b^{k-4} + \cdots + a_2) + \textcolor{red}{a_1}) + \textcolor{blue}{a_0} \\&= \cdots\end{aligned}$$



# Base- $b$ Expansions

$$\begin{aligned}n &= a_k b^k + a_{k-1} b^{k-1} + a_{k-2} b^{k-2} + \cdots + a_2 b^2 + a_1 b + a_0 \\&= b(a_k b^{k-1} + a_{k-1} b^{k-2} + a_{k-2} b^{k-3} + \cdots + a_2 b + a_1) + \textcolor{red}{a_0} \\&= b(b(a_k b^{k-2} + a_{k-1} b^{k-3} + a_{k-2} b^{k-4} + \cdots + a_2) + \textcolor{red}{a_1}) + \textcolor{blue}{a_0} \\&= \cdots\end{aligned}$$

To construct the base- $b$  expansion of an integer  $n$ ,

- Divide  $n$  by  $b$  to obtain  $\textcolor{blue}{n = bq_0 + a_0}$ , with  $0 \leq a_0 < b$
- The remainder  $a_0$  is the rightmost digit in the base- $b$  expansion of  $n$ . Then divide  $q_0$  by  $b$  to get  $\textcolor{blue}{q_0 = bq_1 + a_1}$  with  $0 \leq a_1 < b$
- $a_1$  is the second digit from the right. Continue by successively dividing the quotients by  $b$  until **the quotient is 0**



# Algorithm: Constructing Base- $b$ Expansions

```
procedure base b expansion( $n, b$ : positive integers with  $b > 1$ )  
   $q := n$   
   $k := 0$   
  while ( $q \neq 0$ )  
     $a_k := q \bmod b$   
     $q := q \operatorname{div} b$   
     $k := k + 1$   
  return( $a_{k-1}, \dots, a_1, a_0$ ) {  $(a_{k-1} \dots a_1 a_0)_b$  is base  $b$  expansion of  $n$  }
```

# Example

- $(12345)_{10} = (30071)_8$



# Example

■  $(12345)_{10} = (30071)_8$

$$12345 = 8 \cdot 1543 + 1$$

$$1543 = 8 \cdot 192 + 7$$

$$192 = 8 \cdot 24 + 0$$

$$24 = 8 \cdot 3 + 0$$

$$3 = 8 \cdot 0 + 3$$



# Binary Addition of Integers

$$a = (a_{n-1}a_{n-2} \dots a_1a_0), \quad b = (b_{n-1}b_{n-2} \dots b_1b_0)$$

**procedure** *add*(*a, b*: positive integers)

{the binary expansions of *a* and *b* are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}

*c* := 0

**for** *j* := 0 to *n* − 1

*d* :=  $\lfloor (a_j + b_j + c)/2 \rfloor$

*s*<sub>*j*</sub> :=  $a_j + b_j + c - 2d$

*c* := *d*

*s*<sub>*n*</sub> := *c*

**return**(*s*<sub>0</sub>, *s*<sub>1</sub>, ..., *s*<sub>*n*</sub>) {the binary expansion of the sum is  $(s_n, s_{n-1}, \dots, s_0)_2$ }



# Binary Addition of Integers

$$a = (a_{n-1}a_{n-2} \dots a_1a_0), \quad b = (b_{n-1}b_{n-2} \dots b_1b_0)$$

```
procedure add(a, b: positive integers)
{the binary expansions of a and b are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
c := 0
for j := 0 to n − 1
    d :=  $\lfloor (a_j + b_j + c) / 2 \rfloor$ 
    sj :=  $a_j + b_j + c - 2d$ 
    c := d
sn := c
return(s0, s1, ..., sn) {the binary expansion of the sum is  $(s_n, s_{n-1}, \dots, s_0)_2$ }
```

$O(n)$  bit additions

# Algorithm: Binary Multiplication of Integers

$$a = (a_{n-1}a_{n-2} \dots a_1a_0)_2, \quad b = (b_{n-1}b_{n-2} \dots b_1b_0)_2$$

$$\begin{aligned} ab &= a(b_02^0 + b_12^1 + \dots + b_{n-1}2^{n-1}) \\ &= a(b_02^0) + a(b_12^1) + \dots + a(b_{n-1}2^{n-1}) \end{aligned}$$

```
procedure multiply(a, b: positive integers)
{the binary expansions of a and b are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
for j := 0 to n - 1
    if  $b_j = 1$  then  $c_j = a$  shifted j places
    else  $c_j := 0$ 
{ $c_0, c_1, \dots, c_{n-1}$  are the partial products}
p := 0
for j := 0 to n - 1
    p := p +  $c_j$ 
return p {p is the value of ab}
```

# Algorithm: Binary Multiplication of Integers

$$a = (a_{n-1}a_{n-2} \dots a_1a_0)_2, \quad b = (b_{n-1}b_{n-2} \dots b_1b_0)_2$$

$$\begin{aligned} ab &= a(b_02^0 + b_12^1 + \dots + b_{n-1}2^{n-1}) \\ &= a(b_02^0) + a(b_12^1) + \dots + a(b_{n-1}2^{n-1}) \end{aligned}$$

```
procedure multiply(a, b: positive integers)
{the binary expansions of a and b are  $(a_{n-1}, a_{n-2}, \dots, a_0)_2$  and  $(b_{n-1}, b_{n-2}, \dots, b_0)_2$ , respectively}
for j := 0 to n - 1
    if  $b_j = 1$  then  $c_j = a$  shifted j places
    else  $c_j := 0$ 
{ $c_0, c_1, \dots, c_{n-1}$  are the partial products}
p := 0
for j := 0 to n - 1
    p := p +  $c_j$ 
return p {p is the value of ab}
```

$O(n^2)$  shifts and  $O(n^2)$  bit additions

# Algorithm: Computing div and mod

```
procedure division algorithm (a: integer, d: positive integer)
  q := 0
  r := |a|
  while r ≥ d
    r := r - d
    q := q + 1
  if a < 0 and r > 0 then
    r := d - r
    q := -(q+1)
  return (q, r) {q = a div d is the quotient, r = a mod d is the
  remainder }
```



# Algorithm: Computing div and mod

```
procedure division algorithm (a: integer, d: positive integer)
  q := 0
  r := |a|
  while r ≥ d
    r := r - d
    q := q + 1
  if a < 0 and r > 0 then
    r := d - r
    q := -(q+1)
  return (q, r) {q = a div d is the quotient, r = a mod d is the
  remainder }
```

$O(q \log a)$  bit operations. But there exist more efficient algorithms with complexity  $O(n^2)$ , where  $n = \max(\log a, \log d)$

# Algorithm: Computing div and mod (cont)

```
■ procedure division2 ( $a, d \in \mathbb{N}, d \geq 1$ )  
  if  $a < d$   
    return  $(q, r) = (0, a)$   
   $(q, r) = \text{division2}(\lfloor a/2 \rfloor, d)$   
   $q = 2q, r = 2r$   
  if  $a$  is odd  
     $r = r + 1$   
  if  $r \geq d$   
     $r = r - d$   
     $q = q + 1$   
  return  $(q, r)$ 
```



# Algorithm: Computing div and mod (cont)

```
■ procedure division2 ( $a, d \in \mathbb{N}, d \geq 1$ )  
  if  $a < d$   
    return  $(q, r) = (0, a)$   
   $(q, r) = \text{division2}(\lfloor a/2 \rfloor, d)$   
   $q = 2q, r = 2r$   
  if  $a$  is odd  
     $r = r + 1$   
  if  $r \geq d$   
     $r = r - d$   
     $q = q + 1$   
  return  $(q, r)$ 
```

$O(\log q \log a)$  bit operations.

# Algorithm: Binary Modular Exponentiation

$$b^n = b^{a_{k-1} \cdot 2^{k-1} + \dots + a_1 \cdot 2 + a_0} = b^{a_{k-1} \cdot 2^{k-1}} \dots b^{a_1 \cdot 2} \cdot b^{a_0}$$

Successively finds  $b \bmod m$ ,  $b^2 \bmod m$ ,  $b^4 \bmod m$ ,  $\dots$ ,  $b^{2^{k-1}} \bmod m$ , and multiplies together the terms  $b^{2^j} \bmod m$  where  $a_j = 1$ .

```
procedure modular_exponentiation( $b$ : integer,  $n = (a_{k-1}a_{k-2}\dots a_1a_0)_2$ ,  $m$ : positive integers)
   $x := 1$ 
   $power := b \bmod m$ 
  for  $i := 0$  to  $k - 1$ 
    if  $a_i = 1$  then  $x := (x \cdot power) \bmod m$ 
     $power := (power \cdot power) \bmod m$ 
  return  $x$  { $x$  equals  $b^n \bmod m$ }
```



# Algorithm: Binary Modular Exponentiation

$$b^n = b^{a_{k-1} \cdot 2^{k-1} + \dots + a_1 \cdot 2 + a_0} = b^{a_{k-1} \cdot 2^{k-1}} \dots b^{a_1 \cdot 2} \cdot b^{a_0}$$

Successively finds  $b \bmod m$ ,  $b^2 \bmod m$ ,  $b^4 \bmod m$ ,  $\dots$ ,  $b^{2^{k-1}} \bmod m$ , and multiplies together the terms  $b^{2^j} \bmod m$  where  $a_j = 1$ .

```
procedure modular_exponentiation( $b$ : integer,  $n = (a_{k-1}a_{k-2}\dots a_1a_0)_2$ ,  $m$ : positive integers)
   $x := 1$ 
   $power := b \bmod m$ 
  for  $i := 0$  to  $k - 1$ 
    if  $a_i = 1$  then  $x := (x \cdot power) \bmod m$ 
     $power := (power \cdot power) \bmod m$ 
  return  $x$  { $x$  equals  $b^n \bmod m$ }
```

$O((\log m)^2 \log n)$  bit operations



# Next Lecture

- number theory, cryptography ...

