

# Solution - Exercise [6]

Introduction to Computer Graphics - B-IT Master Course

[Vitaly Kurin]

[Valentin Belonogov]

[Asif Mayilli]

December 15, 2015

## First Exercise

$L_{ambient} = L_a \cdot k_a$ , where

$L_a$  - Intensity of ambient light

$k_a$  - ambient reflection constant, the ratio of reflection of the ambient term present in all points in the scene rendered

$L_{ambient}$  component does not depend on the angle of incidence of the light beam and the view angle of the observer. Ambient light source represents a fixed intensity and fixed color light that affects all objects in the scene equally.

## Second Exercise

$L_{diff} = L_d \cdot r_d \cdot (\mathbf{N} \cdot \mathbf{L})$

$L_{spec} = L_s \cdot r_s \cdot (\mathbf{E} \cdot \mathbf{R})^m$

$N$  - normal on the surface at the point where the light touches the surface

$E$  - the vector in the direction of the viewer

$L$  - the direction vector from the point on the surface towards the light source

$R$  - the vector of a perfectly reflected ray of light ( $L$ )

$m$  - shininess coefficient of the material (the more the surface is mirror-like, the more the coefficient is)

$r_d$  - diffuse reflection constant - the ratio of the reflection of the diffuse term of incoming light

$r_s$  - specular reflection constant specific to the surface - the ratio of the reflection of the specular term of incoming light

$L_d$  - intensity of diffuse term of the light

$L_s$  - intensity of specular term of the light

$L_d$  and  $L_s$  are intensities that are often defined as RGB values. In physics light intensity is measured in Watt per meter squared.