

SWOT High-Rate Raster Data Reveals Antarctic Ice Shelf Motion and Change in Response to Ocean and Ice Dynamics

S. L. Howard, Z. S. Katz, L. Padman, M. R. Siegfried, E. Abrahams, T. M. Snow

Main Data Sources:

SWOT HR Raster 100m data:

- Version C SWOT_L2_HR_Raster_100m at PO.DAAC: <https://doi.org/10.5067/SWOT-RASTER-2.0>
- Version D WOT_L2_HR_Raster_100m at PO.DAAC: <https://doi.org/10.5067/SWOT-RASTER-D>

ICESat-2 ATL06 r006 data

- <https://doi.org/10.5067/ATLAS/ATL06.006>

Corrections:

DAC is available from AVISO through FTP :

- <https://www.aviso.altimetry.fr/en/data/products/auxiliary-products/dynamic-atmospheric-correction.html> <https://doi.org/10.24400/527896/a01-2022.001>

Tides:

To obtain tidal corrections, users must download

- the CATS2008_v2023.nc model from USAP-DC (Howard et al., 2019): <https://doi.org/10.15784/601772>
- TMD 3.0 toolbox from mathworks: <https://www.mathworks.com/matlabcentral/fileexchange/133417-tide-model-driver-tmd-version-3-0> or github : <https://github.com/chadagreene/Tide-Model-Driver>

Additional data:

SWOT HR masks: https://podaac.jpl.nasa.gov/SWOT-events/SWOT_events.html

MEaSURES InSAR-Based Antarctica Ice Velocity Map, Version 2 :

<https://doi.org/10.5067/D7GK8F5J8M8R>

MODIS Mosaic of Antarctica 2003-2004 (MOA2004) at NSIDC :

<https://doi.org/10.5067/68TBT0CGJSOJ>

Ice and tide masks for ICESat-2 ATL14/15 data products on Zenodo:

<https://doi.org/10.5281/zenodo.15327619>