

## 1.1 Difference between deterministic and stochastic world

	deterministic world	stochastic world
Single variable: Temp of a sick man	$R$ $T = 39^\circ C$	random variable $E, Var, \dots$
Variables changing over time: $T$ in first 3 days	$R_+ \rightarrow R$ $T(1) = 39$ $T(2) = 38.5$ $T(3) = 38$ $\vdots$	stochastic process

## 1.2 Difference between various fields of stochastics


Stochastics

- probability theory
- mathematical statistics
- stochastic processes

Consider a pond that contains fish

Prob theory: # of fish at some given time ( $N$ )  
 $E, Var, \text{ or limit laws}$

Mathematical Stats:

Repeat  $m_1, m_2, \dots, m_g$ 

$$(\log \text{ likelihood}) \sum_{k=1}^g P\{\# \text{ marked} = m_k\} \rightarrow \max_N \quad (MLE)$$

1.3 Probability space  $(\Omega, \mathcal{F}, P)$ 

General theory	Bernoulli Scheme $\begin{bmatrix} 1, \text{success} \\ 0, \text{failure} \end{bmatrix}$ $(a_1, \dots, a_n), a_i \in \{0, 1\}$	$[0, 1]$ Select point from
$\Omega$ -sample space	$\#\Omega = 2^n$ , set of all vectors with components $\in \{0, 1\}$	$\Omega = [0, 1]$
$\mathcal{F}$ - $\sigma$ -algebra 1) $\Omega \in \mathcal{F}$ 2) $A \in \mathcal{F} \Rightarrow \Omega \setminus A \in \mathcal{F}$ 3) $A_1, \dots, A_n, \dots \in \mathcal{F}$ $\Downarrow$ $\bigcup_{i=1}^{\infty} A_i \in \mathcal{F}$	$\mathcal{F}$ = power set $\#\mathcal{F} = 2^{\#\Omega} = 2^{2^n}$	$P\{x \in [\alpha, \beta]\}$ $\Rightarrow [\alpha, \beta), (\alpha, \beta],$ $(\alpha, \beta), [\alpha, \beta), \{\beta\} \in \mathcal{F}$ Borel $\sigma$ -algebra
$P$ -probability measure 1) $P(\Omega) = 1$ 2) $A_1, A_2, \dots \in \mathcal{F}$ (disjoint) $\Rightarrow P\{\bigcup_i A_i\} = \sum_i P(A_i)$ $P: \mathcal{F} \rightarrow [0, 1]$	$P\{1\} = p$ $P\{0\} = 1 - p$	$P\{[\alpha, \beta]\} = \beta - \alpha$

1.4 Definition of a stochastic function. Types of stochastic functions.  
 $(\Omega, \mathcal{F}, P)$ Random variable - measurable function  $\xi: \Omega \rightarrow \mathbb{R}$ .

$$\forall B \in \mathcal{B}(\mathbb{R}) : \xi^{-1}(B) \subset \mathcal{F}$$

T - time

 $X: T \times \Omega \rightarrow \mathbb{R}$  - random function, if  $\forall t \in T: X(t, \cdot)$  is a random variable on  $(\Omega, \mathcal{F}, P)$ , denoted  $X_t$

If  $T = \mathbb{R}_+$ , this is called a random process or stochastic process

$T = \mathbb{R}_+^n$ , random field or stochastic field

$T = \mathbb{N}$ , discrete time stochastic process  
or  $\mathbb{Z}$

$T = \mathbb{R}_+ \text{ or } \mathbb{R}$ , continuous time stochastic process


### 1.5 Trajectories and finite-dimensional distributions

$$X: T \times \Omega \rightarrow \mathbb{R}, \quad T = \mathbb{R}_+$$

$\forall t \in T: X_t = X(t, \cdot)$  is a r.v. on  $(\Omega, \mathcal{F}, P)$

Trajectory (= path)

$X_t$  fix  $\omega$  and get mapping  $T \rightarrow \mathbb{R}$




Finite-dimensional distribution  $(X_{t_1}, X_{t_2}, \dots, X_{t_n}), t_1, \dots, t_n \in \mathbb{R}$

In mathematic stats,  $X_{t_1}, X_{t_2}, \dots, X_{t_n}$  are independent

In stochastic process,  $(X_{t_1}, X_{t_2}, \dots, X_{t_n})$  are dependent

Ex:  $X_t = \xi t$ ,  $\xi = \begin{cases} 1, & \text{w.p. } 1/2 \\ 2, & \text{w.p. } 1/2 \end{cases}$



$$P\{X_{t_1} \leq x_1, X_{t_2} \leq x_2\} = \begin{cases} 0, & \min(\frac{x_1}{t_1}, \frac{x_2}{t_2}) < 1 \\ 1/2, & \text{if } \in [1, 2] \\ 1, & \text{if } \geq 2 \end{cases}$$


## 1.6 Renewal process. Counting process.

Renewal processes (discrete time)

$$S_0 = 0, S_n = S_{n-1} + \xi_n, \text{ where } \xi_1, \xi_2, \dots - \text{i.i.d.} > 0 \text{ a.s.}$$

$$P\{\xi_i > 0\} = 1 \Leftrightarrow F(0) = 0$$

$$N_t = \arg \max_k \{S_k \leq t\} \quad (\text{Counting process})$$



$$\{S_n > t\} = \{N_t < n\}$$

$$F \rightarrow \mathbb{E} N_t$$

$$S_n = \xi_1 + \dots + \xi_n$$

## 1.7. Convolution

Convolution  $X \perp\!\!\!\perp Y$ 

$$X \sim F_X, Y \sim F_Y$$

$$F_{X+Y}(x) = \int_{\mathbb{R}} F_X(x-y) dF_Y(y) =: F_X * F_Y$$

conv in terms of distribution functions

$$X \sim p_X, Y \sim p_Y$$

(If  $Y, X$  have densities)

$$p_{X+Y}(x) = \int_{\mathbb{R}} p_X(x-y) p_Y(y) dy =: p_X * p_Y$$

conv in terms of densities

$$S_n = \xi_1 + \dots + \xi_n$$

$$\text{let } F^{n*} := \underbrace{F * \dots * F}_n$$

$$1) F^{n*}(x) \leq F^n(x) \text{ if } F(0)=0$$

$$\xi_1, \dots, \xi_n \stackrel{\text{i.i.d.}}{\sim} F$$

$$\{\xi_1 + \dots + \xi_n \leq x\} \subset \{\xi_1 \leq x, \dots, \xi_n \leq x\} \quad \text{Since } \xi_i \geq 0 \text{ a.s.}$$

$$P\{\xi_1 + \dots + \xi_n \leq x\} \leq \prod_{k=1}^n P\{\xi_k \leq x\}$$

$$\stackrel{||}{F^{n*}(x)} \qquad \qquad \qquad F(x)$$

$$2) F^{n*}(x) \geq F^{(n+1)*}(x)$$

$$\{\xi_1 + \dots + \xi_n \leq x\} \supset \{\xi_1 + \dots + \xi_{n+1} \leq x\}$$

Theorem:  $S_n = S_{n-1} + \xi_n$  where  $\xi_1, \xi_2, \dots \stackrel{\text{i.i.d.}}{\sim} F, F(0)=0$

$$(1) \boxed{U(t) = \sum_{n=1}^{\infty} F^{n*}(t) < \infty}$$

$$(2) \boxed{\mathbb{E}N_t = U(t)}$$

proof for (2)

$$\begin{aligned} \mathbb{E}N_t &= \mathbb{E}[\#\{n: S_n \leq t\}] \\ &= \mathbb{E}\left[\sum_{n=1}^{\infty} \mathbb{1}_{\{S_n \leq t\}}\right] = \sum_{n=1}^{\infty} P\{S_n \leq t\} \\ &= \sum_{n=1}^{\infty} F^{n*}(t) \end{aligned}$$

1.8 Laplace transform. Calculation of an expectation of a counting process (1)

Laplace transform

$$f: \mathbb{R}_+ \rightarrow \mathbb{R} : \mathcal{L}_f(s) = \int_0^{\infty} e^{-sx} f(x) dx$$

$$1) f \text{-density of } \xi, \text{ then } \mathcal{L}_f(s) = \mathbb{E}[e^{-s\xi}]$$

$$2) f_1, f_2 : \mathcal{L}_{\underbrace{f_1 * f_2}_{\text{densities}}}(s) = \mathcal{L}_{f_1}(s) \cdot \mathcal{L}_{f_2}(s)$$

$$3) F \text{-distribution function, } F(0)=0, \quad p = F'$$

$$\mathcal{L}_F(s) = \frac{\mathcal{L}_p(s)}{s}$$



$$\begin{aligned} \text{l.h.s.} &= \int_{\mathbb{R}_+} F(x) \frac{d(e^{-sx})}{s} = - \frac{F(x)e^{-sx}}{s} \Big|_0^{\infty} + \frac{1}{s} \int_{\mathbb{R}_+} p(x) e^{-sx} dx \\ &= \text{r.h.s.} \end{aligned}$$

Ex 1)

$$\begin{aligned} 1) \mathcal{L}_{x^k}(s) &= \int_{\mathbb{R}_+} x^k \frac{d(e^{-sx})}{s} = \frac{n}{s} \int_{\mathbb{R}_+} x^{n-1} e^{-sx} dx \\ &= \frac{n}{s} \cdot \frac{n-1}{s} \cdots \frac{2}{s} \int_{\mathbb{R}_+} e^{-sx} dx = \frac{n!}{s^n} \end{aligned}$$

$$2) \mathcal{L}_{e^{ax}}(s) = \frac{1}{s-a}, \text{ if } a < s$$

1.9 Laplace transform. Calculation of an expectation of a counting process (2)

$$F \rightarrow \mathbb{E}N_t$$

$$\mathbb{E}N_t = U(t) = \sum_{n=1}^{\infty} F^{n*}(t) = F(t) + \left( \sum_{n=1}^{\infty} F^{n*}(t) \right) * F(t)$$

$$\Leftrightarrow U = F + U * F = F + U * p \quad \text{if } F' = p \text{ exists}$$

$\downarrow$  dist. fun.                       $\downarrow$  densities

$$\int_{\mathbb{R}} U(x-y) dF(y) = \int_{\mathbb{R}} U(x-y) p(y) dy$$

$$\mathcal{L}_U(s) = \mathcal{L}_F(s) + \mathcal{L}_U(s) \mathcal{L}_p(s)$$

$$\mathcal{L}_p(s)$$

$$\boxed{\mathcal{L}_U(s) = \frac{\mathcal{L}_p(s)}{s(1 - \mathcal{L}_p(s))}}$$

$$\textcircled{1} F \rightarrow \mathcal{L}_p$$

$$\textcircled{2} \mathcal{L}_p \rightarrow \mathcal{L}_U$$

$$\textcircled{3} \mathcal{L}_U \rightarrow U$$

1.10 Laplace transform. Calculation of an expectation of a counting process (3)

Example:  $S_n = S_{n-1} + \xi_n$ ,  $\xi_1, \xi_2, \dots$  have density  $p(x)$

$$p(x) = \frac{e^{-x}}{2} + e^{-2x}, \quad x > 0$$

$$\mathbb{E}N_t = ?$$

$$\begin{aligned} \textcircled{1} p \rightarrow \mathcal{L}_p : \mathcal{L}_p(s) &= \frac{1}{2} \mathcal{L}_{e^{-x}}(s) + \mathcal{L}_{e^{-2x}}(s) \\ &= \frac{1}{2(s+1)} + \frac{1}{s+2} = \frac{3s+4}{2(s+1)(s+2)} \end{aligned}$$

$$\textcircled{2} \mathcal{L}_p \rightarrow \mathcal{L}_u : \mathcal{L}_u(s) = \frac{\mathcal{L}_p(s)}{s(1-\mathcal{L}_p(s))} = \frac{3s+4}{s^2(2s+3)}$$

$$\begin{aligned} \textcircled{3} \mathcal{L}_u \rightarrow u : \mathcal{L}_u(s) &= \frac{A}{s^2} + \frac{B}{s} + \frac{C}{2s+3} \\ &= \frac{A(2s+3) + B(2s^2+3s) + Cs^2}{s^2(2s+3)} \end{aligned}$$

$$3s+4 = (2B+C)s^2 + (2A+3B)s + 3A$$

$$A = \frac{4}{3}, \quad 2A+3B = 3 \Leftrightarrow B = \frac{1}{9}, \quad 2B+C = 0 \Leftrightarrow C = -\frac{2}{9}$$

$$u(t) = \frac{4}{3}t + \frac{1}{9}(1) - \frac{1}{9}e^{-3/2 t}$$

1.11 Limit theorems for renewal processes

$$S_n = S_{n-1} + \xi_n; \quad \xi_1, \xi_2, \dots \text{ iid } > 0 \text{ a.s.}$$

$$\text{Thm 1 } \mu = \mathbb{E}\xi_1 < \infty \Rightarrow \frac{N_t}{t} \xrightarrow[t \rightarrow \infty]{} \frac{1}{\mu} \text{ a.s.}$$

(Analog to SLLN)

$$\text{SLLN: } \frac{\xi_1 + \dots + \xi_n}{n} \xrightarrow[n \rightarrow \infty]{} \mu \text{ a.s.}$$


Thm 2: (Analog of CLT)  $\sigma^2 = \text{Var } \xi_1 < \infty$

$$\text{Then } Z_t = \frac{N_t - t/\mu}{\sigma \sqrt{t}/\mu^{3/2}} \xrightarrow[t \rightarrow \infty]{} N(0,1)$$

$$P\{Z_t \leq x\} \rightarrow \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-u^2/2} du$$

$$\text{CLT: } \frac{\xi_1 + \dots + \xi_n - n\mu}{\sigma\sqrt{n}} \xrightarrow[n \rightarrow \infty]{} N(0,1)$$

proof (thm 1)



$$\frac{N_t}{S_{N_t+1}} \leq \frac{N_t}{t} \leq \frac{N_t}{S_{N_t}}$$

$$\lim_{t \rightarrow \infty} \frac{N_t}{S_{N_t}} = \lim_{n \rightarrow \infty} \frac{n}{S_n} = \frac{1}{\mu} \text{ by SLLN}$$

$$\lim_{t \rightarrow \infty} \frac{N_t}{S_{N_t+1}} = \lim_{t \rightarrow \infty} \frac{N_t}{N_{t+1}} \cdot \lim_{t \rightarrow \infty} \frac{N_{t+1}}{S_{N_t+1}} = \frac{1}{\mu}$$

$\parallel$   $\parallel$   
 $1$   $1/\mu$

proof (thm 2)

$$P\left\{\frac{S_n - n\mu}{\sigma\sqrt{n}} \leq x\right\} \rightarrow \Phi(x), x \in \mathbb{R}$$

$$P\{S_n \leq n\mu + \sigma\sqrt{n}x\} \rightarrow \Phi(x)$$

$$\Leftrightarrow P\{N_t \geq \frac{t}{\mu}\} \quad \text{(set complements)}$$

$$N\mu \approx t \quad n \approx t/\mu \quad (\text{for } n \text{ large enough})$$

$$n = \frac{t}{\mu} - \frac{\sigma\sqrt{n}}{\mu}x \approx \frac{t}{n} - \frac{\sigma\sqrt{t}}{\mu^{3/2}}x$$

$$\Rightarrow P\{Z_t \geq -x\} \rightarrow \Phi(x) \quad \Leftrightarrow P\{Z_t \leq x\} = 1 - P\{Z_t \geq -x\} \rightarrow 1 - \Phi(-x) = \Phi(x)$$



## Poisson Processes

2.1 Definition of a Poisson process as a special example of a renewal process. Exact forms of the distributions of the renewal process and the counting process (1)


Renewal process

$$S_0 = 0, S_n = S_{n-1} + \xi_n, \xi_1, \xi_2, \dots \text{ i.i.d } > 0 \text{ a.s.}, \xi_i \sim F$$

$$N_t = \arg \max_k \{S_k \leq t\} \quad (\text{Counting process})$$

$$U(t) = \mathbb{E}N_t = \sum_{n=1}^{\infty} F^{*n}(t)$$

$$L_u(s) = \frac{L_p(s)}{s(1-L_p(s))} : p \rightarrow L_p \rightarrow L_u \rightarrow u \quad (p = F')$$



$$L_u(s) = \int_{\mathbb{R}_+} e^{-sx} U(x) dx$$

2.2 ... (2)

Poisson process

Def 1: A process is a renewal process s.t.

$$\xi_i \sim p(x) = \lambda e^{-\lambda x} \mathbb{I}\{x > 0\}, \lambda - \text{intensity or rate}$$

Thm (i): A distribution function of  $S_n$

$$F_{S_n}(x) = \begin{cases} 1 - e^{-\lambda x} \sum_{k=0}^{n-1} \frac{(\lambda x)^k}{k!}, & x > 0 \\ 0, & x < 0 \end{cases}$$

$$p_{S_n}(x) = \lambda \frac{(\lambda x)^{n-1}}{(n-1)!} e^{-\lambda x} \mathbb{I}\{x > 0\}$$

$$(ii) \mathbb{P}\{N_t = n\} = e^{-\lambda t} \frac{(\lambda t)^n}{n!}, N_t \sim \text{Poisson}(\lambda t)$$

2.3 ... (3)

Proof (i)

$$n=1: S_1 = \xi_1$$

$$p_{S_1}(x) = \lambda e^{-\lambda x}, x > 0$$

 $n \rightarrow n+1$ 

$$p_{S_{n+1}}(x) = \int_0^x p_{S_n}(x-y) p_{\xi_{n+1}}(y) dy$$

$$= \int_0^x \frac{\lambda^n (x-y)^{n-1}}{(n-1)!} e^{-\lambda(x-y)} \lambda e^{-\lambda y} dy$$

$$= \frac{\lambda^{n+1}}{(n-1)!} e^{-\lambda x} \int_0^x (x-y)^{n-1} dy = \frac{\lambda^{n+1}}{(n-1)!} e^{-\lambda x} \frac{x^n}{n}$$

$$= \lambda \frac{(\lambda x)^n}{n!} e^{-\lambda x}$$

□

2.4 ... (4)

proof (ii)

$$P\{N_t = n\} = P\{S_n \leq t\} - P\{S_{n+1} \leq t\} \quad (=)$$

$$\{N_t = n\} = \underbrace{\{S_n \leq t\}}_A \cap \underbrace{\{S_{n+1} > t\}}_B$$

$$A \cap B = A \setminus B^c \quad \Rightarrow P\{A \cap B\} = P\{A\} - P\{B^c\}$$

Here:  $B^c \subset A$ 

$$\begin{aligned} &= \left(1 - e^{-\lambda t} \sum_{k=0}^{n-1} \frac{(\lambda t)^k}{k!}\right) - \left(1 - e^{-\lambda t} \sum_{k=0}^n \frac{(\lambda t)^k}{k!}\right) \\ &= e^{-\lambda t} \frac{(\lambda t)^n}{n!} \quad \square \end{aligned}$$

2.5 Memoryless property

A r.v.  $X$  possesses the memoryless property iff

$$P\{X > u+v\} = P\{X > u\} P\{X > v\}. \quad \text{If } P\{X > v\} > 0, \text{ then}$$

$$\boxed{P\{X > u+v \mid X > v\} = P\{X > u\}}$$

Thm 2: Let  $X$  be a r.v. with density  $p(x)$ , then  
 $X$ -memoryless  $\Leftrightarrow p(x) = \lambda e^{-\lambda x}$

Ex buses arrive every  $20 \pm 2$  minutes

$$v = 19 \text{ min}, u = 10 \text{ min}$$

$$(l.h.s.) P\{X > 29 | X > 19\} = 0 \text{ given the data}$$

$$(r.h.s.) P\{X > 10\} = 1$$

Thus, Poisson process is not appropriate

## 2.6. Other definitions of Poisson processes (1)

Def 2  $N_t$  - an integer value process s.t.

$$0) N_0 = 0 \text{ a.s.}$$

$$1) N_t \text{ has independent increments: } \forall t_0 < t_1 < \dots < t_n, \\ N_{t_1} - N_{t_0}, \dots, N_{t_n} - N_{t_{n-1}} \text{ are independent}$$

$$2) N_t \text{ has stationary increments} \\ N_t - N_s \stackrel{d}{=} N_{t-s}$$

$$3) N_t - N_s \sim \text{Poisson}(\lambda(t-s)), t > s$$

$$3) \Rightarrow 2)$$

## 2.7 Other definitions of Poisson processes (2)

$$P\{N_{t+h} - N_t = 0\} = 1 - \lambda h + o(h), h \rightarrow 0$$

$$P\{N_{t+h} - N_t = 1\} = \lambda h + o(h), h \rightarrow 0$$

$$P\{N_{t+h} - N_t \geq 2\} = o(h), h \rightarrow 0$$

$$\lim_{h \rightarrow 0} \frac{1 - P\{N_{t+h} - N_t = 0\}}{h} = \lim_{h \rightarrow 0} \frac{1 - e^{-\lambda h}}{h} = \lambda$$

Def 3  $N_t$  is a Poisson process, if

$$0) N_0 = 0$$

$$1) N_t \text{ has independent increments}$$

$$2) N_t \text{ has stationary increments}$$

$$3') \lim_{h \rightarrow 0} \frac{P\{N_{t+h} - N_t \geq 2\}}{P\{N_{t+h} - N_t = 1\}} = 0$$

2.8 Non-homogeneous Poisson processes (1)

$$N_t \sim \text{Pois}(\lambda t) \Rightarrow \mathbb{E}N_t = \lambda t$$

Def: Let  $\Lambda(t)$  be a differentiable, increasing function s.t.  $\Lambda(0)=0$ . Then,  $X_t = N_t$  is a non-homogeneous Poisson process if,

if, 0)  $N_0 = 0$

1)  $N_t$  has independent increments

2)  $N_t - N_s \sim \text{Pois}(\Lambda(t) - \Lambda(s))$

2.9 Non-homogeneous Poisson processes (2) (NHPP)

$\lambda(t) = \Lambda'(t)$  - intensity function

Properties NHPP:

1)  $\mathbb{E}N_t = \Lambda(t)$

$\Lambda(t) = \alpha t^\beta, \alpha > 0, \beta > 0$  (for example)

2) if  $\lambda(t) = \text{const} \Rightarrow \Lambda(t) = \text{const} \cdot t$

3)  $\Lambda(t)$  - differentiable  $\Rightarrow \Lambda(t)$  - continuous  
 $\Lambda(t)$  - increasing  $\Rightarrow$

$\Rightarrow \exists \Lambda^{-1}(t)$ . If Image  $\Lambda(t) = \mathbb{R}_+$ ,  $N_{\Lambda^{-1}(t)}$  - homogeneous P.P.

2.10 Relation between renewal theory and NHPP (1)

$$S_n = \argmin_t \{N_t = n\}, \quad \xi_n = S_n - S_{n-1}$$

$\xi_1, \xi_2, \dots$  - i.i.d.?

1)  $p_{\xi}(x) = \lambda(x) e^{-\Lambda(x)}$

$$P\{\xi_1 \leq x\} = P\{S_1 \leq x\} = P\{N_x \geq 1\} = 1 - P\{N_x = 0\} \quad \textcircled{=}$$

$$\{S_n > t\} = \{N_t < n\}$$

$\textcircled{=} 1 - e^{-\Lambda(x)}$

Take derivatives of both sides to finish proof  $\square$ .



2.11 ... (2)

$$S_k = \argmin_t \{N_t = k\}$$

$$\xi_k = S_k - S_{k-1}$$

$$1) p_{\xi_1}(t) = \lambda(t) e^{-\Lambda(t)}$$

$$2) p_{\xi_2|\xi_1}(t|s) = \lambda(t+s) e^{-\Lambda(t+s) + \Lambda(s)}$$

$$F_{(\xi_1, \xi_2)}(s, t) = P\{\xi_1 \leq s, \xi_2 \leq t\} = \int_0^s P\{\xi_1 \leq s, \xi_2 \leq t \mid \xi_1 = y\} p_{\xi_1}(y) dy$$

Since  $y \leq s$

$$= \int_0^s P\{N_{t+y} - N_y \geq 1 \mid \xi_1 = y\} p_{\xi_1}(y) dy$$

independent

$$= \int_0^s (1 - e^{-\Lambda(t+y) + \Lambda(y)}) \lambda(y) e^{-\Lambda(y)} dy$$

$$p_{(\xi_1, \xi_2)}(s, t) = \frac{\partial}{\partial t} \left( \frac{\partial}{\partial s} F_{(\xi_1, \xi_2)}(s, t) \right)$$

$$= \frac{\partial}{\partial t} \left( (1 - e^{-\Lambda(t+s) + \Lambda(s)}) \lambda(s) e^{-\Lambda(s)} \right)$$

$$= \lambda(t+s) e^{-\Lambda(t+s) + \Lambda(s)} \lambda(s) e^{-\Lambda(s)}$$

Then  $p_{\xi_2|\xi_1}(t|s) = \frac{p_{(\xi_1, \xi_2)}(s, t)}{p_{\xi_1}(s)}$  finishes the proof  $\square$ .

2.12 ... (3)

$\xi_1, \xi_2, \dots$  - i.i.d. ? (NHPP can be obtained from renewal process iff NHPP is homogeneous PP)

$$p_{\xi_1}(t) = p_{\xi_2|\xi_1}(t|s), \quad \forall t, s > 0$$

$$\lambda(t) e^{-\Lambda(t)} = \lambda(t+s) e^{-\Lambda(t+s) + \Lambda(s)}$$

$$\left( \int_0^T \dots dt \right) : e^{-\Lambda(0)} - e^{-\Lambda(T)} = e^{-\Lambda(s)} - e^{-\Lambda(T+s) + \Lambda(s)}$$

$$\Lambda(T) = \Lambda(T+s) - \Lambda(s), \quad \forall s, T > 0$$

$\Lambda(t)$  - increasing

$$\Rightarrow \Lambda(t) = \lambda t$$

const



2.13 Elements of queuing theory.  $M/G/k$  systems (1)

$$\begin{aligned}
 P\{N_{t+h} - N_t = 0\} &= 1 - \lambda h + o(h) \\
 P\{N_{t+h} - N_t = 1\} &= \lambda h + o(h) \\
 P\{N_{t+h} - N_t \geq 2\} &= o(h)
 \end{aligned}$$

 $M/G/k$ 

I) Arrival Process:  $M$  - memoryless (Poisson)  
 $D$  - deterministic  
 $G$  - general

II) Service time ( $M, D, G$ )III) A number of services ( $1, 2, \dots, \infty$ ) $M/G/\infty$  $\tau > 0$  (time moment)

$N(t)$   
 Customer arrivals  $\rightarrow$   $N_1(t)$  - still being served at  $\tau : \lambda_1(t) = \lambda(1 - G(\tau - t))$   
 $\rightarrow$   $N_2(t)$  - already completed by  $\tau : \lambda_2(t) = \lambda G(\tau - t)$

$$\begin{aligned}
 P\{N_1(t+\delta) - N_1(t) = 1\} &= P\{N(t+\delta) - N(t) = 1\} \cdot (P\{Y > \tau - t\} + o(\delta)) \\
 &= (\delta\lambda + o(\delta)) (1 - G(\tau - t) + o(\delta)) \\
 &= \boxed{\lambda\delta(1 - G(\tau - t) + o(\delta))}
 \end{aligned}$$

## 2.14 ... (2)

$$P\{N_1(t) = n_1, N_2(t) = n_2\} = P\{N_1(t) = n_1, N_2(t) = n_2 \mid N(t) = n_1 + n_2\} \cdot P\{N(t) = n_1 + n_2\}$$

$$= C_{n_1+n_2}^{n_1} (1 - G(\tau - t))^{n_1} G(\tau - t)^{n_2} \cdot e^{-\lambda t} \frac{(\lambda t)^{n_1+n_2}}{(n_1+n_2)!}$$

$$= \frac{\lambda t (1 - G(\tau - t))^{n_1}}{n_1!} e^{-\lambda(1 - G(\tau - t))} \cdot \frac{\lambda t (G(\tau - t))^{n_2}}{n_2!} e^{-\lambda G(\tau - t)}$$

$$= P\{N_1(t) = n_1\} \cdot P\{N_2(t) = n_2\}$$

Therefore  $N_1 \perp N_2$

## 2.15 Compound Poisson Processes (1)

$$X_t = \sum_{k=1}^{N_t} \xi_k, \quad \xi_1, \xi_2, \dots \text{ i.i.d. }, N_t \text{ - P.P. with intensity } \lambda$$

and  $\xi_1, \xi_2, \dots$  and  $N_t$  are independent

$\xi_1, \xi_2, \dots$  claim sizes

$N_t$  - amount of claims until time  $t$  (Insurance interpretation)

$X_t$  - aggregated claim amount

### 1) Probability generating function (PGF)

$\xi$  - integer,  $\geq 0$  values

$$\boxed{\phi_\xi(u) = \mathbb{E}[u^\xi], \quad |u| < 1}$$

$$\xi_1 \perp \xi_2 \Rightarrow \phi_{\xi_1 + \xi_2}(u) = \phi_{\xi_1}(u) \phi_{\xi_2}(u)$$

### 2) Moment-generating function (MGF)

$$\boxed{L_\xi(u) = \mathbb{E}[e^{-u\xi}], \quad \xi \geq 0, u > 0}$$

## 2.16 ... (2)

### 3) Characteristic function

$$\phi_\xi(u) = \mathbb{E}[e^{iu\xi}], u \in \mathbb{R}, \forall \xi, \quad \phi_\xi: \mathbb{R} \rightarrow \mathbb{C}, \quad \xi_1 \perp \xi_2 \Rightarrow \phi_{\xi_1 + \xi_2}(u) = \phi_{\xi_1}(u) \phi_{\xi_2}(u)$$

$$\text{Thm } \boxed{\phi_{X_t - X_s}(u) = e^{\lambda(t-s)(\phi_\xi(u) - 1)}}$$

$$\begin{aligned} \text{Proof: } \text{lhs} &= \mathbb{E} e^{iu(X_t - X_s)} = \sum_{k=0}^{\infty} \mathbb{E} \left[ e^{iu(X_t - X_s)} \mid N_t - N_s = k \right] P\{N_t - N_s = k\} \\ &= \sum_{k=0}^{\infty} (\phi_\xi(u))^k e^{-\lambda(t-s)} \frac{[\lambda(t-s)]^k}{k!} \end{aligned}$$

$\xi_1 + \dots + \xi_k$  from 11

□

2.17 ... (3)

$$X_t = \sum_{k=1}^{N_t} \xi_k, \quad \xi \text{ can be any random variable}$$

$$\xi: \phi_\xi(u) = E[e^{iu\xi}]$$

$$\phi: \mathbb{R} \rightarrow \mathbb{C}$$

$$\xi_1 \perp \xi_2 \Rightarrow \phi_{\xi_1 + \xi_2}(u) = \phi_{\xi_1}(u) \phi_{\xi_2}(u)$$

$$\text{Thm: } \phi_{X_t - X_s}(u) = e^{\lambda(t-s)(\phi_\xi(u) - 1)}, \quad t > s \geq 0$$

proof:

$$\text{lhs} = E[e^{iu(X_t - X_s)}]$$

$$= \sum_{k=0}^{\infty} E[e^{iu(X_t - X_s)} | N_t - N_s = k] \cdot P\{N_t - N_s = k\}$$

$\downarrow$  since  $\xi_1, \xi_2, \dots \perp N_t \sim \text{Pois}(\lambda(t-s))$   
 $\xi_1, \dots, \xi_k$

$$= \sum_{k=0}^{\infty} [\phi_\xi(u)]^k \cdot e^{-\lambda(t-s)} \frac{[\lambda(t-s)]^k}{k!} \quad \square$$

2.18 ... (4)

$$\text{Corollary: } \begin{cases} EX_t = \lambda t E\xi, \\ \text{Var } X_t = \lambda t E\xi^2 \end{cases}$$

proof:  $E[\xi^r] < \infty \Rightarrow \phi(u)$  is  $r$ -times differentiable at 0  
 and  $\phi^{(r)}(0) = i^r E\xi^r$

$$EX_t = \frac{\phi'_{X_t}(0)}{i} = \frac{\lambda t (\phi'_\xi(0) \cdot \phi_{X_t}(0))}{i} = \lambda t E\xi, \quad \square$$

$\downarrow$   
 $i = E\xi$