

Lagrange Multipliers

Lagrange Multipliers

Equality Constraint

Inequality Constraint

Lagrange multipliers, a.k.a. undetermined multipliers, are used to find **the stationary points** of a function of several variables subject to one or more constraints.

Equality Constraint

Sometime it may be difficult to find a analytic solution of the constraint between variables x_i . We can introduce a parameter λ called Lagrange multiplier. From a geometrical perspective, consider a D -dimensional variable $\mathbf{x} = (x_1, \dots, x_D)^T$. The constraint equation $g(\mathbf{x}) = 0$ is a $(D - 1)$ -dimensional surface in \mathbf{x} -space.


Lemma-1. At any point on the surface $g(\mathbf{x}) = 0$ the gradient ∇g will be orthogonal to the surface.

Proof. Consider a point \mathbf{x} on the surface, and consider a nearby point $\mathbf{x} + \epsilon$ that also on the surface. If we make a Taylor expansion around this point, we have:

$$g(\mathbf{x} + \epsilon) \approx g(\mathbf{x}) + \epsilon^T \nabla g(\mathbf{x}) \quad (1)$$

because $g(\mathbf{x} + \epsilon) = g(\mathbf{x})$, we have $\epsilon^T \nabla g(\mathbf{x}) \approx 0$. And if $\|\epsilon\| \rightarrow 0$, we have $\epsilon^T \nabla g(\mathbf{x}) = 0$. And because at this time ϵ is parallel to the surface $g(\mathbf{x}) = 0$, so ∇g is normal to the surface.

A geometrical picture of the technique of Lagrange multipliers in which we seek to maximize a function $f(\mathbf{x})$, subject to the constraint $g(\mathbf{x}) = 0$. If \mathbf{x} is D dimensional, the constraint $g(\mathbf{x}) = 0$ corresponds to a subspace of dimensionality $D - 1$, indicated by the red curve. The problem can be solved by optimizing the Lagrangian function $L(\mathbf{x}, \lambda) = f(\mathbf{x}) + \lambda g(\mathbf{x})$.



Lemma-2. If \mathbf{x}^* on $g(\mathbf{x}) = 0$ and $\forall \mathbf{x}$ on $g(\mathbf{x}) = 0$, we have $f(\mathbf{x}^*) \geq f(\mathbf{x})$, then $\nabla f(\mathbf{x}^*)$ is orthogonal to the surface $g(\mathbf{x}) = 0$.

Proof. Otherwise we could increase the value of $f(\mathbf{x})$ by moving a short distance along the surface.

From *Lemma-1* and *Lemma-2* we have:

$$\nabla f + \lambda \nabla g = 0 \quad (2)$$

where $\lambda \neq 0$ is the Lagrange multiplier.

Then we can define the *Lagrangian* function

$$L(\mathbf{x}, \lambda) := f(\mathbf{x}) + \lambda g(\mathbf{x}) \quad (3)$$

And

$$\begin{aligned} \nabla_{\mathbf{x}} L &= 0 \Rightarrow \nabla f + \lambda \nabla g = 0 \\ \frac{\partial L}{\partial \lambda} &= 0 \Rightarrow g(\mathbf{x}) = 0 \end{aligned}$$

Thus to find the maximum of f s.t. $g(\mathbf{x}) = 0$, we solve the equation


$$\begin{cases} \nabla_{\mathbf{x}} L = 0 & \Rightarrow \nabla f + \lambda \nabla g = 0 \\ \frac{\partial L}{\partial \lambda} = 0 & \Rightarrow g(\mathbf{x}) = 0 \end{cases} \quad (4)$$

and get a $(D + 1)$ -dimension equations to get stationary points.

Inequality Constraint

The problem of maximizing f with inequality constraint of the form $g(\mathbf{x}) \geq 0$.

Illustration of the problem of maximizing $f(\mathbf{x})$ subject to the inequality constraint $g(\mathbf{x}) \geq 0$.



There are two kinds of solution possible.

- *inactive* constraint: The stationary point lies in the region $g(\mathbf{x}) > 0$. The function $g(\mathbf{x})$ is useless, $\lambda = 0$.
- *active* constraint: The stationary point lies in the surface $g(\mathbf{x}) = 0$. This is analogous to the equality constraint case, $\lambda \neq 0$.

Note that now the sign of λ is crucial, because f only be at a maximum if its gradient is oriented away from the region $g(\mathbf{x}) > 0$. Therefore $\lambda > 0$ (if we want a minimum, we set $\lambda < 0$).

For either cases, we have $\lambda g(\mathbf{x}) = 0$. Then we can define:

KKT(Karush-Kuhn-Tucker) conditions

$$\begin{aligned} g(\mathbf{x}) &\geq 0 \\ \lambda &\geq 0 \\ \lambda g(\mathbf{x}) &= 0 \end{aligned}$$

And the solution is also straightforward.