

Discrete Structures Section 2

KP. Wang

Assignment #3

Solutions

1. Section 1.7 # 14, 18, 19, 20, 25, 28
2. Section 1.8 # 2, 7, 17
3. Section 2.1 # 10, 20, 21, 25, 32, 44
4. Section 2.2 # 14, 15
5. Consider sets A, B , and C , show that $A \times (B \cap C) = (A \times B) \cap (A \times C)$ by mutual subsets.
 1. *Solution:* If x is rational and not zero, then by definition we can write $x = p/q$, where p and q are nonzero integers. Since $1/x$ is then q/p and $p \neq 0$, we can conclude that $1/x$ is rational.
 2. *Solution:*
 - (a) We must prove the contrapositive: If n is odd, then $3n + 2$ is odd. Assume that n is odd. Then we can write $n = 2k + 1$ for some integer k . Then $3n + 2 = 3(2k + 1) + 2 = 6k + 5 = 2(3k + 2) + 1$. Thus $3n + 2$ is two times some integer plus 1, so it is odd.
 - (b) Suppose that $3n + 2$ is even and that n is odd. Since $3n + 2$ is even, so is $3n$. If we add subtract an odd number from an even number, we get an odd number, so $3n - n = 2n$ is odd. But this is obviously not true. Therefore our supposition was wrong, and the proof by contradiction is complete.
3. *Solution:* The proposition we are trying to prove is "If 0 is a positive integer greater than 1, then $0^2 > 0$." Our proof is a vacuous one. Since the hypothesis is false, the conditional statement is automatically true.
4. *Solution:* We need to prove the proposition "If 1 is a positive integer, then $1^2 \geq 1$." The conclusion is the true statement $1 \geq 1$. Therefore the conditional statement is true. This is an example of a trivial proof, since we merely showed that the conclusion was true.
5. *Solution:* Suppose by way of contradiction that a/b is a rational root, where a and b are integers and this fraction is in lowest terms (that is, a and b have no common divisor greater than 1). Plug this proposed root into the equation to obtain $a^3/b^3 + a/b + 1 = 0$. Multiply through by b^3 to obtain $a^3 + ab^2 + b^3 = 0$. If a and b are both odd, then the left-hand side is the sum of three odd numbers and therefore must be odd. If a is odd and b is even, then the left-hand side is odd + even + even, which is again odd. Similarly, if a is even and b is odd, then the left-hand side is even + even + odd, which is again odd. Because the fraction a/b is in simplest terms, it cannot happen that both a and b are even. Thus in all cases, the left-hand side is odd, and therefore cannot equal 0. This contradiction shows that no such root exists.
6. *Solution:* There are two things to prove. For the "if" part, there are two cases. If $m = n$, then of course $m^2 = n^2$; if $m = -n$, then $m^2 = (-n)^2 = (-1)^2 n^2 = n^2$. For the "only if" part, we suppose that $m^2 = n^2$. Putting everything on the left and factoring, we have $(m + n)(m - n) = 0$. Now the only way that a product of two numbers can be zero is if

one of them is zero. Therefore we conclude that either $m + n = 0$ (in which case $m = -n$), or else $m - n = 0$ (in which case $m = n$), and our proof is complete.

7. *Solution:* The cubes that might go into the sum are 1, 8, 27, 64, 125, 216, 343, 512, and 729. We must show that no two of these sum to a number on this list. If we try the 45 combinations $(1 + 1, 1 + 8, \dots, 1 + 729, 8 + 8, 8 + 27, \dots, 8 + 729, \dots, 729 + 729)$, we see that none of them works. Having exhausted the possibilities, we conclude that no cube less than 1000 is the sum of two cubes.
8. *Solution:* There are several cases to consider. If x and y are both nonnegative, then $|x| + |y| = x + y = |x + y|$. Similarly, if both are negative, then $|x| + |y| = (-x) + (-y) = -(x + y) = |x + y|$, since $x + y$ is negative in this case. The complication (and strict inequality) comes if one of the variables is nonnegative and the other is negative. By the symmetry of the roles of x and y here (strictly speaking, by the commutativity of addition), we can assume without loss of generality that it is x that is nonnegative and y that is negative. So we have $x \geq 0$ and $y < 0$.
9. *Solution:* The equation $|a - c| = |b - c|$ is equivalent to the disjunction of two equations: $a - c = b - c$ or $a - c = -b + c$. The first of these is equivalent to $a = b$, which contradicts the assumptions made in this problem, so the original equation is equivalent to $a - c = -b + c$. By adding $b + c$ to both sides and dividing by 2, we see that this equation is equivalent to $c = (a + b)/2$. Thus there is a unique solution. Furthermore, this c is an integer, because the sum of the odd integers a and b is even.
10. *Solution:*
 - a) true,
 - b) true,
 - c) false,
 - d) true,
 - e) true - the one element in the set on the left is an element of the set on the right, and the sets are not equal,
 - f) true,
 - g) false - the two sets are equal.
11. *Solution:*
 - a) The empty set has no elements, so its cardinality is 0.
 - b) This set has one element (the empty set), so its cardinality is 1.
 - c) This set has two elements, so its cardinality is 2.
 - d) This set has three elements, so its cardinality is 3.
12. *Solution:*
 - a) $\{\emptyset, \{a\}\}$
 - b) $\{\emptyset, \{a\}, \{b\}, \{a, b\}\}$
 - c) $\{\emptyset, \{\emptyset\}, \{\{\emptyset\}\}, \{\emptyset, \{\emptyset\}\}\}$

13. *Solution:* We need to prove two things, because this is an "if and only if" statement. First, let us prove the "if" part. We are given that $A \subseteq B$. We want to prove that the power set of A is a subset of the power set of B , which means that if $C \subseteq A$ then $C \subseteq B$. But this follows directly from Exercise 17. For the "only if" part, we are given that the power set of A is a subset of the power set of B . We must show that every element of A is also an element of B . So suppose a is an arbitrary element of A . Then $\{a\}$ is a subset of A , so it is an element of the power set of A . Since the power set of A is a subset of the power set of B , it follows that $\{a\}$ is an element of the power set of B , which means that $\{a\}$ is a subset of B . But this means that the element of $\{a\}$, namely a , is an element of B , as desired.
14. *Solution:* In each case the answer is a set of 3 -tuples.
- a) $\{(a, x, 0), (a, x, 1), (a, y, 0), (a, y, 1), (b, x, 0), (b, x, 1), (b, y, 0), (b, y, 1), (c, x, 0), (c, x, 1), (c, y, 0), (c, y, 1)\}$
- b) $\{(0, x, a), (0, x, b), (0, x, c), (0, y, a), (0, y, b), (0, y, c), (1, x, a), (1, x, b), (1, x, c), (1, y, a), (1, y, b), (1, y, c)\}$
- c) $\{(0, a, x), (0, a, y), (0, b, x), (0, b, y), (0, c, x), (0, c, y), (1, a, x), (1, a, y), (1, b, x), (1, b, y), (1, c, x), (1, c, y)\}$
- d) $\{(x, x, x), (x, x, y), (x, y, x), (x, y, y), (y, x, x), (y, x, y), (y, y, x), (y, y, y)\}$
15. *Solution:* In each case we want the set of all values of x in the domain (the set of integers) that satisfy the given equation or inequality.
- a) It is exactly the positive integers that satisfy this inequality. Therefore the truth set is $\{x \in \mathbb{Z} \mid x^3 \geq 1\} = \{x \in \mathbb{Z} \mid x \geq 1\} = \{1, 2, 3, \dots\}$.
- b) The square roots of 2 are not integers, so the truth set is the empty set, \emptyset .
- c) Negative integers certainly satisfy this inequality, as do all positive integers greater than 1. However, $0 \not\geq 0^2$ and $1 \not\geq 1^2$. Thus the truth set is $\{x \in \mathbb{Z} \mid x < x^2\} = \{x \in \mathbb{Z} \mid x \neq 0 \wedge x \neq 1\} = \{\dots, -3, -2, -1, 2, 3, \dots\}$.
16. *Solution:* Since $A = (A - B) \cup (A \cap B)$, we conclude that $A = \{1, 5, 7, 8\} \cup \{3, 6, 9\} = \{1, 3, 5, 6, 7, 8, 9\}$. Similarly $B = (B - A) \cup (A \cap B) = \{2, 10\} \cup \{3, 6, 9\} = \{2, 3, 6, 9, 10\}$.
17. *Solution:* a) This proof is similar to the proof of the dual property. Suppose $x \in \overline{A \cup B}$. Then $x \notin A \cup B$, which means that x is in neither A nor B . In other words, $x \notin A$ and $x \notin B$. This is equivalent to saying that $x \in \bar{A}$ and $x \in \bar{B}$. Therefore $x \in \bar{A} \cap \bar{B}$, as desired. Conversely, if $x \in \bar{A} \cap \bar{B}$, then $x \in \bar{A}$ and $x \in \bar{B}$. This means $x \notin A$ and $x \notin B$, so x cannot be in the union of A and B . Since $x \notin A \cup B$, we conclude that $x \in \overline{A \cup B}$, as desired.
- b) The following membership table gives the desired equality, since columns four and seven are identical.

A	B	$A \cup B$	$\overline{A \cup B}$	\overline{A}	\overline{B}	$\overline{A} \cap \overline{B}$
1	1	1	0	0	0	0
1	0	1	0	0	1	0
0	1	1	0	1	0	0
0	0	0	1	1	1	1

18. *Solution:* First we will show that $A \times (B \cap C) \subseteq (A \times B) \cap (A \times C)$. Suppose $(a, b) \in A \times (B \cap C)$. By definition of the Cartesian product, this means $a \in A$ and $b \in B \cap C$. By definition of intersection, it follows that $b \in B$ and $b \in C$. Thus, since $a \in A$ and $b \in B$, it follows that $(a, b) \in A \times B$. Also, since $a \in A$ and $b \in C$, it follows that $(a, b) \in A \times C$. Now we have $(a, b) \in A \times B$ and $(a, b) \in A \times C$, so $(a, b) \in (A \times B) \cap (A \times C)$. We've shown that $(a, b) \in A \times (B \cap C)$ implies $(a, b) \in (A \times B) \cap (A \times C)$ so we have $A \times (B \cap C) \subseteq (A \times B) \cap (A \times C)$.

Next we will show that $(A \times B) \cap (A \times C) \subseteq A \times (B \cap C)$. Suppose $(a, b) \in (A \times B) \cap (A \times C)$.

By definition of intersection, this means $(a, b) \in A \times B$ and $(a, b) \in A \times C$. By definition of the Cartesian product, $(a, b) \in A \times B$ means $a \in A$ and $b \in B$. By definition of the Cartesian product, $(a, b) \in A \times C$ means $a \in A$ and $b \in C$. We now have $b \in B$ and $b \in C$, so $b \in B \cap C$, by definition of intersection. Thus we've deduced that $a \in A$ and $b \in B \cap C$, so $(a, b) \in A \times (B \cap C)$. In summary, we've shown that $(a, b) \in (A \times B) \cap (A \times C)$ implies $(a, b) \in A \times (B \cap C)$ so we have $(A \times B) \cap (A \times C) \subseteq A \times (B \cap C)$.

The previous two paragraphs show that $A \times (B \cap C) \subseteq (A \times B) \cap (A \times C)$ and $(A \times B) \cap (A \times C) \subseteq A \times (B \cap C)$, so it follows that $(A \times B) \cap (A \times C) = A \times (B \cap C)$.