

Regenerative host • fixed physiology

Finite ring replacement | 2 tapes × 4 initial bodies × 2 deletion layouts | 256 steps


A Saved density fields | input tape 202610195601




t = 0 (given ring)



greedy, t = 128



greedy, t = 256



uniform, t = 256

Density rho: 0–0.5983
Purple → yellow
Same spatial crop

A prescribed ring persists;
its density redistributes.

Not de novo ring formation.

B All 16 trajectories | joint window test

Initial body	5601 G	5601 U	5602 G	5602 U
Compact	FAIL	FAIL	FAIL	FAIL
Ring	PASS	PASS	PASS	PASS
Double lobe	FAIL	FAIL	FAIL	FAIL
Separated	FAIL	FAIL	FAIL	FAIL

G = original prestate greedy; U = uniform.
PASS: all four 64-step windows; last-two result identical.

C Live birth-cohort mass at t = 256

Tape	Layout	Founder	Born>0	Born>128
5601	Greedy	36.87	35.69	22.03
5601	Uniform	36.60	35.66	22.07
5602	Greedy	36.18	35.31	22.00
5602	Uniform	35.98	35.43	21.95

Founder initially 64; Born>128 is a subset of Born>0.
Both labels are observers, never inputs to the rule.

Only ring passes (4/16); original greedy already supports renewal.
Renewal is supported here; cue learning was tested in a different, non-renewing host.