

# A Contact recedes before the second resource site opens

Compact h2 · acquisition: A only → shared state 512 → B future (retain shown)





State 512 → B future
Retain / reset / shared
Actor increments off




ρ: 0–1 M/cell · linear


Fixed window: x6–29, y2–25 (y up); 24 × 24 cells. Strips show site boundaries.




B contact is already present during acquisition.
Its decline precedes availability.
Test access: 16 of every 64 actions;
stock outside the available site is inaccessible.

# B At B, weak uptake accompanies a large tissue difference


Same compact h2 receiver · B available only in four 16-action windows




Both start with E = 3.221 EU; retain/reset end at 0.3445 / 0.0029 EU.
More tissue can persist with weak intake; U and motor accumulate from action 513.

# C All eight pairs: more energy, less motor payment

Four acquired histories × two sites · endpoint 768 · retain minus reset





Retain − reset (all four quantities)
Shared − reset (E only): all positive; spatial benefit varies

A: more final contact, but less M in three pairs. Compact h2/A: less U despite more E.

# D Earlier receivers: the energy effect reverses with access schedule

Continuous → continuous
Continuous → intermittent
Intermittent → continuous
Intermittent → intermittent




Acquisition → test · separate 512–640

ΔE (EU) · shared marks in orange · all tested futures lose tissue