
Inherited tolerance and injury experience have different consequences


Formation 0–128 (T00) → Histories 129–160 → Test 161–224

N: no prior cut; C: 10% +x cuts before 129 / 145. N and C use paid P/F transport (T11).

Test: none or 20% +x cut before 161. T00: no prior cut, no transport capability or transport charge.

A Compact offspring survives cuts that kill its parent





Light: parent; dark: offspring


Compact R × I follow-up (N/x, C/x)

	I0	I1
R0	178 / 183: zero	178 / 184: zero
R1	both >0 at 224	both >0 at 224

R1I0 C − N, x: ΔM +.687; ΔE −6.279; ΔT −40.633 (T = full-cycle body-time).

Density ρ: white 0 → teal 1 M/cell. Same x28–67, y28–63 crop; y up.


B Positive interaction does not make injury experience beneficial



Multiring: all six matched child − parent M / E / body-time < 0

R/I follow-up: At each I, R1 − R0: M / E / T < 0 in all four matched paths. R × I is nonzero.


Parent and child: all six original futures grow at every final-32 update.




Fixed 68 × 64 crop; same ρ scale; smaller than A.

Multiring: interaction +6.298, but both C − N totals < 0.

C Regaining total mass is not restoration of the missing structure





Compact: extinct 273.
Multiring: old total M regained at 312; cut region ends at 22.78% of control.

Cut region peaks at 325, then falls. Shape L1 = 373.060 after registration (displacement 0): not restored.

Separate state 256 excision. Fixed x10–85, y10–85 crop.

Grey: uninjured region; orange: cut region. This is a different receiver and clock from A/B.

One compound mutation per parent; one tape / +x scenario. Learning off; no stable-recovery or general-adaptation claim.